

2004/2005

Forest Genetics Council of BC

Tree Improvement Program Project Report

FGC

Front Cover

Research collection of the douglas-fir cone worm, *Dioryctria abietivorella*, reared from infested spruce and douglas-fir cones.

Adult females of the spruce seed chalcid, *Megastigmus piceae*. The one on the right is ovipositing into an ovule.

Sporulating lesion of White Pine Blister Rust, *Endocronartium ribicola*, on a white pine branch.

Forest Genetics Council of BC

Tree Improvement Program Project Report

Coordinated and compiled by:
Beacon Hill Communications Group Inc.

and

Roger Painter
Tree Improvement Branch
British Columbia Ministry of Forests

Acknowledgements

It is my pleasure to present this report for the eighth year. The Tree Improvement Program has proven to be one of the most enduring and successful partnerships sponsored by our provincial government. What started out as a short-term funding source to augment current budgets and increase annual production has expanded into a complex, progressive, and successful public-private partnership that encompasses all aspects of tree improvement. This is now been recognized by other areas within the provincial government. It was nominated by the Ministry of Forests for the Premier's award and was one of three programs reviewed under the Auditor General's report on Managing for Results. In the past year we have again added yet another sub-program to target a broad need within tree improvement. The newest effort is specifically geared to answer problems about seed pest management by investing in longer-term research and problem solving in that vital area. We look forward to results from seed pest research to provide us with more valuable tools for protecting our seed orchards and crops.

The Forest Investment Council continues to recognize the value of investing in tree improvement. There is little doubt that this support is providing us with the resources to continue with a vibrant and robust long-term program. We continue to build on the management processes of the program. These approaches, including business planning and performance management and long-term strategic planning, are key to the success of the program. Other organizations in addition to the Forest Genetics Council (FGC) view them as a successful model for developing investment programs on the co-operative partnerships model.

I would like to acknowledge the project leaders for their contributions to this report. This publication continues to provide a well-deserved display of our work, and the program receives numerous compliments on the successes of tree improvement from various people in the forest industry. My congratulations to you all for an excellent eighth annual report. I hope you enjoy it. My sincere thanks for the hard work of our editorial review team for taking the time to look over our submissions. They include Diane Douglas, Ward Strong, Ron Planden, and Greg O'Neil.

Our sincere thanks to Dr. Robb Bennett and the Seed Pest Management Program for their valuable contributions to the front and back covers of this year's report. Seed Pest Management has expanded in the past year and now is looking forward to an increased presence in providing a strong research element in their work.

Again, my thanks to all those who have worked on this program in the past year, including the review committees, species committees and various Technical Advisory Committees (TACs). You are the reason that this program has a strong technical direction. To all the project leaders, I hope 2005/2006 is as successful as the previous year.

Roger A. Painter
Tree Improvement Co-ordinator
Forest Genetics Council

Forest Genetics Council Co-chairs

Once again we thank the many industry and government staff who contribute to the provincial gene resource management program for their cooperation and dedication over the past year. Although forestry in BC is changing quickly, there is a consistent and solid base of support for tree improvement, and a well-established planning process in place.

During the 2004/05 fiscal year, substantial progress was made in several important areas:

- Chief Forester's Standards for Seed Use, in support of the new *Forest and Range Practices Act*, were completed. The Standards consolidate existing policy, guidelines, and practices and result in efficiencies for both industry and government stakeholders. Many members of council and affiliated committees contributed to and improved the Standards during the past year.
- Council membership changes were completed, allowing long-serving councillors a rest and new energy to be brought to bear.
- Under the direction of the recently appointed Chief Forester Jim Snetsinger, the FIA – Tree Improvement Program continues to support the long-term goals of Council for increased seed quality and seed production in seed orchards.

The coming year will continue to present additional challenges for Council. As programs mature and industry continues to feel pressure to increase competitiveness in global commodity markets, the need to realize value from all silviculture investments will increase. Seed pests in our British Columbia seed orchards took a significant toll on interior species and will be the focal point of a new Forest Genetics Council sub-program. Climate change is a growing challenge to forest management policy development, and we can expect that seed selection and use will need to be reviewed in this regard in the short term. This will allow us to meet the public expectations regarding stewardship of the genetic resource on public lands. Your program Co-chairs are confident that Council and its affiliated committees will rise to meet these challenges in 2005/06.

On a final note, this year marks the final contribution of Shane Browne-Clayton to the Forest Genetics Council as Industry Co-Chair. Shane has worked tirelessly and with consistent co-operation since the onset of the newly structured council in 1996. As Ministry Co-Chair, I extend my thanks and that of countless others for the contribution and friendship offered by Shane over this period.

Shane Browne-Clayton
Industry Co-chair

Dale Draper
Ministry of Forests
Co-chair

The Forest Genetics Council Co-Chairs invite you to review the programs and projects described in this report and return any questions or comments to:

TIP Program Administrator
Ministry of Forests
Tree Improvement Branch
PO Box 9518 Stn Prov Govt
Victoria, B. C. V8W 9C2

Further Tree Improvement information can be found at our Web sites:

Forest Genetics Council	http://www.fgcouncil.bc.ca
Ministry of Forests, Tree Improvement Branch	http://www.for.gov.bc.ca/TIP/index.htm
Ministry of Forests, Research Branch	http://www.for.gov.bc.ca/research/

FGC

Table of Contents

Forest Genetics Council of BC Tree Improvement Program Project Report i
Acknowledgements ii
Forest Genetics Council Co-chairs iii
Tree Improvement in British Columbia iv
Ministry of Forests Program Overview iv
Forest Genetics Council of B.C. iv
FIA – Tree Improvement Subprograms v
1.0 Expansion of Orchard Seed Supply Subprogram (SelectSeed Company Ltd.) 1
2.0 Gene Resource Information Management 2
3.0 Centre for Forest Gene Conservation (CFGC) 3
4.0 Tree Breeding 5
4.1 Coastal Douglas-fir 5
4.2 Sitka Spruce 5
4.3 Western Hemlock and True Fir Forest Genetics Programs 6
4.4 Western Redcedar Breeding Program 7
4.5 Yellow-cedar Breeding Program 8
4.6 Coastal White Pine 8
4.7 Interior Douglas-fir 10
4.8 Interior Spruce 11
4.9 Genetic Resistance for Western Gall Rust (*Endocronartium harknessii*) in Lodgepole Pine 12
4.10 Western White Pine Blister Rust Resistance Screening: Ongoing in the Interior 14
4.11 Ponderosa Pine Reselection Work at the Skimikin Seed Orchard Site 14
4.12 Western Larch 16
5.0 Operational Tree Improvement – The Eighth Year in Review 17
5.1 Orchard Projects 18
5.1.1 Saanich Forestry Centre 18
5.1.2 Sechelt Seed Orchard 21
5.1.3 Mount Newton Seed Orchard 21
5.1.4 Bowser Seed Orchards 22
5.1.5 Kalamalka Seed Orchards 23
5.1.6 Vernon Seed Orchard Company (VSOC) 25
5.1.7 Grandview Seed Orchards 26
5.1.8 Tolko Industries – Eagle Rock Seed Orchards 27
5.1.9 Prince George Tree Improvement Station (PGTIS) 27
5.1.10 Skimikin Seed Orchard 29
5.1.11 Saanich Seed Orchards 31
5.2 Technical Support Projects 32
5.2.1 Cone and Seed Pest Management – Interior Operations 32

5.2.2 Western Redcedar Pollen Monitoring, Saanich Peninsula and Sechelt (SPU0212) 32
5.2.3 Damage Potential of Field Densities of *Leptoglossus occidentalis* 35
5.2.4 Douglas-fir and Western Redcedar Cone and Seed Insect Pest Monitoring Services to Coastal Conifer Seed Orchard Managers 37
5.2.5 Estimation of Deleterious Effects of Different Levels of Inbreeding on Western Hemlock Seed Orchard Production 38
5.2.6 Determination of Selfing Rates in a Top-pruned Interior Spruce Seed Orchard (SPU 410) 39
5.2.7 Development of Pollen Management Guidelines for Yellow-cedar 39
5.2.8 Improving Seed Production in Yellow-cedar Seed and Breeding Orchards 41
5.2.9 Operational Crown Management Trials in Various Orchards: Interior Spruce High-density Seed Orchard #600 and Interior Douglas-fir Orchard #304 43
5.2.10 Testing the Effects of Upgraded Irrigation Delivery Systems in NE Mid-interior Spruce Seed Orchard #305 and Interior Lodgepole Pine Seed Orchard #307 44
5.2.11 Clonal Verification of a Spruce Sheerit Orchard (#172) Using DNA Markers (SPU 0611) 45
5.2.12 Enhancing Seed Production in North Okanagan Lodgepole Pine Seed Orchards (SPU 0710) 46
5.2.13 Chemical Control Techniques Against *Dioryctria abietivorella* in Interior Fd and Pw Orchards 48
"Where Do Work" Photo Collage 50
6.0 Extension and Communications 52
6.1 Extension Technical Advisory Committee (ETAC Subprogram) 52
7.0 Seed Pest Management 54
7.1 Operational Assay Program for Fungi Attacking Conifer Seedlin BC 54
7.2 Identification of an Effective Sex Pheromone Lure for the Fir Coneworm, *Dioryctria abietivorella*, and Demonstration of Its Efficacy in Seed Orchards (SPR 0002) 54
7.3 Identification of Seed Chalcids Infesting Seeds of BC Conifers (SPR 0003) 54
7.4 Identification of Communication Signals Produced by Male Western Conifer Seed Bugs, *Leptoglossus occidentalis* (SPR 0001) 55
Appendix 1 PGC Seed Planning Units 56
Appendix 2 Tree Species 57
Appendix 3 Discussions by Tree Species 58
Appendix 4 Authors and Species Topic 59
Appendix 5 Author Contact List 63

List of Figures

Figure 1. Relationship between FGC Strategic Plan, FGN-TIP, and participants in various forest gene resource management areas 4
Figure 2. Schematic of organizational relationships among FGC, FGN, SelectSeed and seed users 5
Figure 3. Summary of SelectSeed orchard developments, and progress in propagation and planting during the year ending March 31, 2005 7
Figure 4. Natural Stand Seed Planning Zones 8
Figure 5. Summary of western redcedar polygenic tests 9
Figure 6. Relationship between cold-related damage and seed source at a high-elevation coastal polygenic test near Harrison Mills 9
Figure 7. Summary of the 15-year results from the Mt. Robson interior Douglas-fir progeny tests (209 families) measured in fall 2004 10
Figure 8. Relationship between predicted parental breeding values for wood relative density (age 14) and individual tree volume at age 40 in Skeena Kitsumkalum interior spruce at the Skeeksis Seed Orchard 11
Figure 9. Summary of results from the Fort Nelson and Nelson interior spruce progeny tests measured in fall 2004 12
Figure 10. Blackwater Creek 118 OP families 13
Figure 11. Blackwater on Tutu (Moon Lady/Living Tree) 110 OP families 13
Figure 12. Blackwater Creek 118 OP Families 14
Figure 13. Summary of the 10-year results from the East Kootenay Series 3 western larch open-pollinated progeny tests (160 families) measured in fall 2004 16
Figure 14. Number of projects and funding by region 17
Figure 15. Number of projects by areas of investment 18
Figure 16. Weighted Average gain for SelectSeed in IBTP orchards only 20
Figure 17. Pollen management activities by project 24
Figure 18. Orchard composition activities by project 24
Figure 19. Break down of projects 25
Figure 20. Mean daily western redcedar pollen catch (pollen grains/mm2) for three stations at Sechelt. 30
Figure 21. Mean daily western redcedar pollen catch (pollen grains/mm2) for five stations on the Saanich peninsula. Black bars include other species of the same size and configuration (predominately red alder). 30
Figure 22. 30
Figure 23. 37
Figure 24. Average inflated scales per conelet counts (SEC) for eight of eleven coastal Douglas-fir seed orchards 37
Figure 25. Propensity to self. Ranking of the clones from the lowest to the highest level of selfing 39
Figure 26. Effects of crown position on seed yield per cone in a top grafted interior spruce seed orchard 39
Figure 27. Relationship between yellow-cedar pollen viability at pollination) and population origin elevation and mean temperature 40
Figure 28. Yellow-cedar pollination study: % filled seed) for four mating types 40
Figure 29. Yellow-cedar pollen development and viability (+/- s.e.) over time (2003 to 2004) 41
Figure 30. Yellow-cedar pollen development and viability (+/- s.e.) over time (2003 to 2004) 42

Figure 15. Yellow cedar pollen development and viability (+/- s.e.) over time (2004 to 2006) 26
Figure 16. Yellow cedar pollen development and viability (+/- s.e.) over time (2004 to 2006) 26
Figure 17. Yellow cedar pollen development and viability (+/- s.e.) over time (2004 to 2006) 26
Figure 18. Yellow cedar pollen development and viability (+/- s.e.) by site over time (2004 to 2006) 26
Figure 19. Treatments applied to seeds in the spruce provenance orchard 27
Figure 20. Treatments applied to seeds in ITE orchard #226 in 2006 27
Figure 21. Ice cream topographical effect on crop 28
Figure 22. Cone verification in Sitka spruce using micro-satellite markers 28
Figure 23. Temperature (T) and vapour pressure deficits (VPD) cone and precipitation (mm) for the period of April 15 to May 15 for the study period of 2000 to 2006 37
Figure 24. Temperature (T) and vapour pressure deficit (VPD) cone and precipitation (mm) for the period of May 16 to May 31 for the study period of 2000 to 2006 37
Figure 25. Mean total seed per cone (TSPC), filled seed per cone (FSPC) and seed weight for cones with (Bag) and without (Nobag) insect protection for the study period of 2000 to 2006 37
Figure 26. Mean cone number per sugar whorl branch (pollen cone clusters, first-year conelets, and second-year cones), and cone weights (first-year conelets and second-year cones) for the study period of 2000 to 2006 38
Figure 27. Orchard block trials. Percentage of cones infested with Boxgalls in Acrolix sprayed with different concentrations of Dimethoate 400E 39
Figure 28. Individual tree trials. Percentage of cones infested with Boxgalls on individual trees sprayed with different concentrations of Dimethoate 400E 40
Figure 29. SP 758 (a) and 1013 trial locations 50

List of Plates

Plates 1 and 2. 1998 parents from Series 4 exhibit frost damage at a high-elevation coastal site near Harrison Mills 9
Plate 3. Barry Jacquish presents Douglas-fir provenance results at the Brendon Seed Orchard to ITA participants 10
Plate 4. Ponderosa pine Resistance Work at the Brendon Seed Orchard Site 11
Plate 5. Western White Pine Blister Rust Resistance Screening 11
Plate 6. Yellow cypress clonal trial selections 12
Plate 7. J. Russell discussing wood quality and breeding at Coastal Silviculture Committee meeting 13
Plate 8. Our first search assistant seedbed three to (13) 21
Plate 9. Cone collection with grafts nearby 22
Plates 10 to 15. Equipment activities at PGTS 23
Plate 16. Trees being removed from old plantations 24
Plate 17. Prepping the site for a spruce provenance trial 24
Plate 18. Spraying in the Bulkley Valley spruce orchards 25
Plate 19. Mike Carlson discusses the results of birch provenance trials at the Brendon Seed Orchard with ITA participants 26
Plate 20. Planting replacement grafts in the rooting bed 26
Plate 21. Elongicta albifrontella adult 28
Plate 22. Elongicta damage on Douglas-fir cones 28
Plates 23 and 24. Photos from the forest genetics tour held October 5, 2006 to view some of the provenance/progeny trials in the Kitwanga/Kispiox area of BC 32

Tree Improvement in British Columbia

Ministry of Forests Program Overview

Forest gene resource management encompasses the conservation, controlled use, and enhancement of genetic resources of forest tree species, and related communication and extension activities. The Forest Genetics Council of British Columbia (FGC) coordinates a provincial forest gene resource management program and its implementation by stakeholders in the forest industry, the Ministry of Forests (MoF), Canadian Forest Service (CFS), and universities.

The Forest Investment Account (FIA) is a major funding agency for forest gene resource management in British Columbia. Through the Tree Improvement Program (TIP), FIA invests in forest gene resource management activities that support the objectives and are consistent with existing government and industry activities.

The FIA-MoF Tree Improvement Program is guided by strategic and annual business plans prepared by the FGC.

Forest Genetics Council of B.C.

The Forest Genetics Council of British Columbia (FGC) is a multi-stakeholder group representing the forest industry, the Ministry of Forests (MoF), Canadian Forest Service (CFS), and universities. Council's mandate is to champion forest gene resource management in British Columbia, to provide strategic and business planning for a co-operative provincial forest gene resource management program, and to advise the province's Chief Forester on forest gene resource management policies. FGC members provide strategic direction to the provincial forest gene resource management program. FGC Technical Working Committees (TWC) provide technical and policy information to Council and contribute to the development of FGC plans and associated budgets.

Council's goal is to maximize the economic benefits from tree improvement gains in wood quality, quantity, and pest resource by supporting a gene conservation program, developing a crop tree production capacity, doubling the average volume gain of select seed, and increasing the amount of select seed used. The FGC Business Plan defines the annual set of activities and associated budgets to achieve this goal.

Forest gene resource management is a co-operative effort. In broad terms, the MoF funds tree breeding activities and private industry funds operational production of reforestation materials. The Canadian Forest Service, MoF Research Branch, and universities undertake research supporting tree improvement, while private institutions focus on applied research related to operational production.

"Select" refers to seed from trees from breeding programs that select those with desirable characteristics for growth, strength, or pest resistance. "Seed" or seed from elite is an advanced reforestation material, including registered orchard seed. The Forest Genetics Council does not support genetic engineering, and no genetically modified materials are used for forest seed reforestation in British Columbia.

[redacted]: Test improvement Subprograms

[redacted]

- [redacted]
- [redacted]
- [redacted]
- [redacted]
- [redacted]
- [redacted]
- [redacted]

[redacted]

[redacted]

[redacted]

[redacted]

[redacted]

[redacted]



Seedplanning Unit						2003 Seedling Planting		2004 Seedling Planting		2005 Seedling Planting		2006 Seedling Planting		
Rgn	SPZ	Elev	SIBEC	Location	Total ssh	ssh planted	SPI done	ssh planted	ssh done	ssh planted	ssh done	ssh planted	ssh done	% complete
Kal	9E	<1000	01	#97	2200	97	97	91	91	345	218	2135	2139	97%
Kal	3J	all	01	#101	975	0	0	0	0	45	0	760	797	81%
Kal	4L	all	01	#101	780	0	0	0	0	80	60	775	748	95%
Kl	9E	<1400	07	#97	1500	0	116	116	116	78	28	955	936	98%
Kl	11J	<1400	08	#97	4798	111	668	777	668	1,413	1,214	4750	4,589	96%
Kl	11L	<1200	09	#101	4500	736	780	736	780	990	814	4605	4,238	94%
Kl	11L	<1200	07	#101	4871	613	638	613	638	641	407	4880	4,695	96%
Sk	13J	<1400	09	Riverside	1815	418	471	418	471	280	237	2480	2,150	89%
Sk	8M	1200	09	#101	1500	227	255	227	255	210	164	2070	1,974	95%
Sk	8M	1200	08	Sunshine	2100	0	0	0	0	0	0	2105	2,100	94%
Sk	11J	>1400	08	#101	2100	116	140	116	140	110	110	2070	2,060	94%
Sk	11F	>1400	07	Sunshine	2000	80	84	80	84	85	35	1980	1,915	96%
Sp	11J	1200-1400		Riverside	1400	91	94	91	91	10	10	1480	1,480	99%
Sp	11J	<1200	101	Riverside	816	21	45	21	21	10	24	805	845	98%
		Total			31,355	3,116	3,774	3,116	3,414	3,847	3,446	34,564	33,315	94%

Figure 1: Summary of Seed Stock orchard developments, and progress in propagation and planting during the year ending March 31, 2005.

2.0 Gene Resource Information Management

Leslie McAuley

Gene resource information management (GRIM) forms a critical link to the long-term stewardship and sustainable resource management of the province's forest tree gene resources over future generations. The key objectives of GRIM are to:

- Develop a provincial GRIM framework to support land use and forest development/stewardship plans, forest regeneration (planting) programs, forest genetics research, and gene resource management (gene conservation/seed deployment) (GRM) strategies.
- Support and maintain GRM registries and data repositories.
- Provide access to tree improvement program products (seed and vegetative material for operational use).
- Build an effective GRM monitoring program integrated with broader forestry (i.e., Criteria & Indicators) and land-based resource management initiatives.
- Develop a means to incorporate genetic gain into timber supply analyses through forest inventory update procedures, forest simulator modelling tools, and the use of GIS-ready spatial and attribute GRM data sets.

GRIM project accomplishments in 2004 /05 included:

- Strategic planning initiated for the development of a Resource Information Strategy for the province's genetic resources, including preliminary development of Gene Resource Management plans (for each data set).
- Development of a web-based parent tree registry and on-line registration functions in support of the new Chief Forester's Standards for Seed Use. Both new features will be accessible through the Seed Planning and Registry system (SPAR).
- Enhancement of genetic resource data sets in support of the new Chief Forester's Standards for Seed Use, including revised Seed Planning Unit data sets and development of new Tested Parent Tree Areas-of-Use data subsets for species plans and seed registration.
- Development of draft resource attribute and mapping criteria to assist in the superior provenance resource

feature (source, quality) review project.

- Development of quick, easy-to-view and printable Look Up theme maps to assist resource managers.
- Enhancement of the web-based SeedMap tool to adopt a standardized look and feel (SOLE, VEG, FOR HEALTH, GENETIC RESOURCES map layer folders) and incorporate new spatially enabled queries (e.g., SEED deployment).
- Development of land-based queries (Source: RESULTS system) to support indicators (Genetic Diversity, Gene Conservation, Forest Productivity) required for national Criteria & Indicators reporting and provincial State of the Forests and Forest Resource Evaluation program reporting.
- Integration of CAIR within broader government-wide strategic planning and forest management initiatives and analyses (Mountain Pine Beetle, Inventory Update, Silviculture, Timber Supply).
- Development of CAIR results-based information reporting requirements under the new *Forest and Range Practices Act* (on-going).
- Development of web-based training modules, tutorials, and on-line Help (on-going).

Figure 4. Natural Stand Seed Planning Zones

3.0 Centre for Forest Gene Conservation (CFGC)

Sally Aitken

The Centre for Forest Gene Conservation at the University of British Columbia continues to work towards the gene conservation objectives of the Forest Genetics Council. Our projects address strategies and theory for gene conservation, evaluate current levels of protection of forest genetic resources, assess the potential impacts and opportunities for mitigating the effects of climate change on forest genetic resources, and investigate species-specific genetic issues associated with at-risk and rare tree species. We also strive to meet the needs for extension in the area of gene conservation. Here we highlight our recent progress and publications in these areas.

Cataloguing and Documenting Protection of Forest Genetic Resources

The evaluation of the current *in situ* conservation status of all tree species in BC through modelling ranges and

species density and the estimation of the representation inside and outside of protected areas are both fairly complete. Two publications have been completed (Hamann et al. 2004, Hamann et al. submitted), and analyses are complete for a third. However, ground-truthing is required for some species in some areas. This will require field work in the upcoming fiscal year. The *ex situ* resources in seed storage at the Tree Seed Centre have been analyzed spatially and will be strategically evaluated along with *in situ* genetic resources.

Climate Change and Gene Conservation

This project area is currently the largest we have underway, and Andreas Hamann, Tongli Wang, and Pia Smets have made enormous progress in the past year (note that this research is partially funded through an NSERC Strategic/BIOCAP Canada grant). We now have an excellent fine-scale climate model for British Columbia (Hamann and Wang 2005). This model has since been developed to incorporate and downscale regional climate-change predictions in collaboration with David Spittlehouse. A software tool called ClimateBC is nearing release that will allow users to predict current and future monthly and annual climatic means for any location in BC The development of these models and tools has been the focus of our climate-change research in 2004-05. In 2005-06, we will be able to move on to further applications of these models. Two important ones are evaluating the future protection of tree species in protected areas given climate change and revising SPU boundaries for seed transfer based on climatic rather than geographic information.

Genetic Structure of Minor Species

1. The PhD research of Andy Bower on whitebark pine genecology, inbreeding depression, and the interaction between inbreeding depression and disease resistance is nearing completion. Whitebark pine has tremendous mid-winter cold hardiness (LT50 below -70°C) and a greater degree of cold hardiness during the active growing season than most conifers (LT50 -9°C). Populations show significant variation in fall, but not spring, cold hardiness, with higher hardiness in more northern populations. This geographic pattern will need to be considered when planning restoration activities for this species. Earlier CFGC research found high levels of inbreeding in two BC populations of whitebark pine. In extending this assessment to provenances from further south, it appears that inbreeding rates decline with latitude. Inbreeding levels in conifers usually decline as stands age. However, there is a tendency for inbreeding levels to increase with age in whitebark pine in stands that are heavily infected with white pine blister rust.
2. A project on genetic diversity and the mating system of Pacific dogwood was initiated in 2004 by Karolyn Keir, who successfully developed six micro-satellite markers for this species from those developed for Florida dogwood. Her preliminary work indicates this species has very low levels of genetic diversity. There is anecdotal evidence from Idaho that low genetic diversity contributed to high mortality due to the disease Anthracnose. She will be returning to start an M.Sc. degree in September, 2005 on this project.
3. Five hundred Garry oak seedlings from 12 families were grown at UBC in the past year for a preliminary look at quantitative variation and to give us experience in the cultivation of this somewhat recalcitrant species. Joanne Tuytel has established a small, common-garden experiment in raised nursery beds this spring. Extra seedlings will be used to establish a Garry oak meadow at the UBC Botanical Garden. If acorn crops are good this year, collections will be made for initiation of a genecological study.

Extension

Our extension activities have grown exponentially in the past year. Requests for information, analyses, and presentations, particularly on the climate-change research, are high, coming from provincial government agencies in BC and Alberta, from the Canadian Forestry Service, and from other countries and organizations. Our website (www.forestry.ubc.ca/cfgc) has summaries of all projects and extensive electronic resources for others. We are nearing completion of a major upgrade in both the structure and the design of the website.

CFGC Publications in 2004-05

Aitken, S.N. 2004. Genecology and adaptation of forest trees. Pp. 197-204 *In*: Jeffrey Burley, Julian Evans, and John A. Youngquist (eds.). Encyclopedia of Forest Sciences. Four-volume set. Elsevier.

Bennuah, S.Y., T. Wang, and S.N. Aitken. 2004. Genetic analysis of the *Picea sitchensis x glauca* introgression zone in British Columbia. For. Ecol. Mgmt. 197(1-3): 65-77.

Gapare, W. J., S.N. Aitken, and C.E. Ritland. 2005. Genetic diversity of core and peripheral Sitka spruce (*Picea sitchensis* (Bong.) Carr) populations: implications for conservation of widespread species. Biological Conservation 123: 113-123.

Hamann A. and T.L. Wang. 2005. Models of climatic normals for genecology and climate change studies in British Columbia. Agricultural and Forest Meteorology 128 (3-4): 211-221

Hamann, A., S. N. Aitken, A. D. Yanchuk. 2004. Cataloguing *in situ* protection of genetic resources for major commercial forest trees in British Columbia. For. Ecol. Mgmt. 197(1-3): 295-305.

Bennuah, S.Y., T. Wang, and S.N. Aitken. 2004. Genetic analysis of the *Picea sitchensis x glauca* introgression zone in British Columbia. For. Ecol. Mgmt. 197(1-3): 65-77.

4.0 Tree Breeding

4.1 Coastal Douglas-fir

Michael Stoehr, Keith Bird, Clint Hollefreund

Coastal Douglas-fir (SPU 01 and SPU 31)
Breeding work in the spring resulted in four breeding-population sublines being fully completed and three sublines being partially completed (general combining ability component only). In total, over 100 full-sib families and over 65 half-sib (general combining ability test) were sown in the fall for out-planting in early spring of 2006. Site maintenance was carried out on 10 progeny test sites and five realized-gain sites. Increment cores, for the determination of wood density in second-generation progeny tests (EP 708) were collected and are being analyzed. This completes the collection of cores from 66 sites. High-elevation tests were analysed and breeding values determined. These were used to rogue CANFOR orchard #116.

Sub-maritime Douglas-fir (SPU 19)
A total of 91 (half-sib) families created from polymixes from orchard #181 were grown for the establishment of three progeny tests in the coastal-interior transition zone. Test sites selected are in the D'Arcy and Pemberton areas at elevations from 500 to 1,100 m. From these tests, a total of 15 parents will be backward selected starting at age 5. Orchard #181 will be supplemented with 10 forward selections of the genecology tests planted in 1996 on four transition-zone sites (EP 1200). The four EP 1200 sites in the transition zone were maintained.

4.2 Sitka Spruce

John King, Dave Ponsford

Weevil resistance program
Last year we established the first phase of the F1 breeding program – weevil resistance. Parents were designated by their phenotypic resistance to the weevil and their source (mainly Haney or Big Qualicum). Some susceptible parents were also used. The trials were established in the Adam's River drainage on north Vancouver Island and at Jordan River, south Vancouver Island. Another site that we held over was established this spring (2005) on Queen Charlotte Island. We sowed a second series of the F1s this spring to be established in spring 2006.

Selection and breeding work continues, but our focus in the future for the breeding program will be to identify parents for specific traits. They will still be field assessed for resistance after weevil augmentation, but a more detailed microscopic evaluation is being undertaken in order to classify parents according to putative resistance mechanisms. This is particularly valuable for "constitutive" type of mechanisms such as sclereid cells or constitutive resin cells, and we are working to extend these techniques to look at some of the "inducible" mechanisms, particularly traumatic resin cell production.

In the past we have relied strongly on the Canadian Forestry Service for entomology back-up for this program. With their new directions we expect to rely more on the universities, although of course Dr. René Alfaro still provides major input.

Ongoing trial assessments for weevil attack have continued, including some of the hazard assessment trials. These were a series of trials established to test resistance over a variety of ecological and potential weevil-hazard areas. The main objective is to aid in determining deployment guidelines. This exercise will be carried out with Don Heppner, the MoF Coast regional entomologist, and we hope to provide regional guidelines for spruce over the next year.

Major articles published in the last year include:

King J.N. and R.I. Alfaro. 2004. Breeding for resistance to a shoot weevil of Sitka spruce in British Columbia, Canada. In: M.J., Carson and C. Walter, eds. Plantation Forest Biotechnology for the 21st Century. Forest Research, New Zealand.

King J.N., R.I Alfaro, and C. Cartwright. 2004. Genetic resistance of Sitka spruce (*Picea sitchensis*) populations to the white pine weevil (*Pissodes strobi*): distribution of resistance. Forestry 77:269-278.

Alfaro R.I., L van Akker, B.Jaquish, and J. King. 2004. Weevil resistance of progeny derived from putatively resistant and susceptible interior spruce parents. For. Ecol. and Mgmt. 202:369-377.

Other Components – QCI, NW Hybrid Program

The Queen Charlotte Island (QCI) program allows us to assess Sitka growth free from weevil attack and to support the Seed Orchard program (SO 142 and its replacement). Our major difficulty has been in getting "free-to-grow" conditions, in this case not from weevil or brush but from deer browse, and a good deal of our maintenance effort is on browse protection. This year we spent a good deal of time cleaning up, tagging, and removing browse protection from the trials we have established there.

The transition program is designed mainly to test seed transferability in a very environmentally heterogeneous area with pressures from both frost and weevil. We completed a five-year assessment this year and had a very successful extension trip showing local foresters in the northwest our program. We have continued some selections of good hybrid weevil-resistant trees for possible orchard deployment.

4.3 Western Hemlock and True Fir Forest Genetics Programs

Charlie Cartwright

Low-elevation Maritime Hemlock (SPU 03)

Hemlock is the most under-utilized, widely used species for regeneration in BC. The breeding program hopes this will be rectified as higher-gain seed becomes available. Since the genetic worth of first-generation orchard seed lots has now levelled at around 16% gain, higher gains can occur only through moving to the next generation of selections. To this point, 220 trees have been chosen from advanced-generation HEMTIC (Hemlock Tree Improvement Co-operative) trials based on age 5 height (the tests incorporate the best parents from first-generation programs in Washington, Oregon, and BC). As age 10 results are tabulated starting in the fall of 2005, about half of these early picks will be culled. It was apparent that the heritability of growth for individual trees was low, as was expected for this species. This means that the value of the selections relies much more on their family performance than on that of individuals within the family. To remedy this for later advanced-generation tests, clonal trials are used. In these, several copies of each individual are planted to allow highly accurate within-family selection. In 2005 three test sites of this type were planted. In the meantime, methods of producing a desirable balance of male and female strobili with young selected material in small pots has commenced in order that breeding with the new material can begin as soon as the age 10 figures are available.

Note: Participation in HEMTIC by the BC Ministry of Forests has been terminated. As per the Memorandum of Understanding with which the co-op was established, age 5 data from the HEMTIC tests have been sent, thus securing access to all HEMTIC-improved material. Continued cooperation with the current members of the co-op is planned. Hopefully further exchanges of data (age 10) can be arranged.

High-elevation Hemlock (SPU 24)

The advantages of high-elevation hemlock include faster growth than yellow cedar and better wood quality than balsam fir. The species plan calls for first-generation testing of about 300 parents to achieve a 15% level of

genetic gain in growth traits. To this point results are available for 240 parents, with new breeding values provided this year for 212 of those parent trees.

All Hemlock SPUs – Seed Transfer Concerns

Further gains from the hemlock breeding program may be achieved through expansion of the planning units. The limits to seed transfer for wild stand seed lots can also be established. At this point it is uncertain how high in elevation and latitude the tested material may be moved. The advanced-generation trials are relatively complex and need good survival to produce useful results. Also, because the environment is quite variable, many small tests are required, whereas in progeny trials a few large tests in good growing conditions produce good results. For this reason, about 35 small hemlock seed transfer tests comprised of wild-stand and tested seed sources have been widely dispersed in BC over the last ten years. Highlights of staff activities this year in these trials include weeding eight of the younger sites and measuring seven. Next year another two test sites will be measured, and then an analysis with the purpose of revising seed transfer boundaries will commence.

Pacific Silver Fir (SPU 09)

Four Pacific silver fir provenance sites were planted in 2002, and eight in 2003. Of the trials, one near Hope was destroyed by wild fire, and a second was inaccessible due to a bridge being out. First data beyond nursery results will be taken on the older series in the 2005/06. As well, replacement planting of the site that burned is planned, along with beefed-up planting of the others.

Grand Fir (SPU 36)

The species subcommittee considered declaring a superior seed source based on age 20 data from eight provenance test sites. The top source (Salmon River) produced 12% more volume than other well-adapted sources for the seed planning unit. Once juvenile/mature correlations were considered, the advantage of this provenance was 8%. The species sub-committee will not recommend the use of this source for B+ seed until it can be ascertained that enough trees remain at the source to make seed collection feasible.

Sub-alpine Fir

Four nursery trials for this species were established in 2002. One had to be abandoned due to poor survival. Of the remaining three, two were in need of intensive maintenance. Measurements for growth on two sites and for Christmas tree traits on one site will be carried out in 2005-06. To cover gene conservation concerns and long-term growth data, excess replacement stock will be out-planted to two new field sites in the coming year. The Balsam Wooly Adelgid quarantine will limit the scope of this, so sowing of new tests targeted at the northern interior of BC will also be undertaken next year.

Noble Fir

As with grand fir, identification of superior provenance to produce B+ seed is under consideration by the species subcommittee. Based on age 16 data from 12 sites, the top source was 11% in volume beyond averages for other well-adapted provenances. After consideration of juvenile/mature correlation, the achievable gain was about 6%. Similar to the situation for grand fir, the viability of using the chosen superior seed source near Mount St. Helen's (French Butte) will be checked in the field. Other good provenances are in ecological reserves/ parks, or on logged sites; the Mt. St Helen's trees are USDA National Forest and young mature.

4.4 Western Redcedar Breeding Program

John Russell, Craig Ferguson

The last series of first-generation polycross testing for the maritime-low SPU have been planted. In total, this project involved the testing of 938 parents across seven annual series and 46 test sites (Figure 5). Breeding values for volume at rotation are currently available for series 1 and 2 based on seven-year heights (288 parents) and series 2, 3, and 4 based on five-year heights (327 parents). Approximately 55 parents from the first four series have been selected for advanced generation breeding and have been established into breeding orchards.

Needle terpene concentrations are currently being evaluated on all 930 parent trees in progeny tests. In addition, sixty within-family selections for high-needle monoterpene concentration/deer resistance were made at the Holt Creek, SVI family/population study. These selections will be bred using a combination of

selfing and outcrossing, and the resultant progeny will be screened for monoterpene concentration and deer resistance. Both backward and forward selections can then be incorporated into a deer-resistant production population.

Heartwood tropolone concentrations have been evaluated for approximately 300 parent trees from the first three series, and the top 50 backward selections have been established in breeding orchards. Tropolone concentration has been shown to be related to wood durability. Further durability screening needs to be developed prior to the operational release of these clones for seed production and further breeding.

			# of test sites/SPU[1]						
Series	Year	#parents	ML	MH	QCI	NM	SM	INT	Total
1	1998	150	3	1	-	-	-	-	4
2	1999	138	4	-	-	-	2	-	6
3	2000	110	3	-	1	1	1	-	6
4	2001	57	4	1	1	-	-	-	6
5	2003	160	3	2	1	1	1	-	8
6	2004	168	2	3	-	1	1	2	9
7	2005	147	4	1	1	-	1	-	7
Total		**930**	**23**	**8**	**4**	**3**	**6**	**2**	**46**

1. ML= Maritime low (under 600 m); MH=Maritime high (>600m); QCI=Queen Charlotte Islands; NM=North Maritime; SM=Submaritime; INT=Interior

Figure 5. Summary of western redcedar polycross tests

Plates 1 and 2. PNW parents from Series 4 exhibit frost damage at a high-elevation coastal site near Harrison Mills



Figure 4. Relationship between cold-related damage and seed source at a high-elevation coastal ponderosa test near Harrison Mills.

4.5 Yellow-cedar Breeding Program

John Russell, Craig Ferguson

There are currently clonal values from approximately 5,000 clones represented in three series of annual testing. Nine-year data from the first series of cloned progeny tests have shown that the top 1% of the clones results in 15% volume gain at 60 years. These new clonal selections are currently being re-propagated for eventual operational release. As new selections come on line from all studies, breeding orchards – both low and high elevation – are being updated. Maintenance continued this year in both field trials and breeding orchards. In addition, the project was engaged in monitoring pollen performance and production for collection for advanced-generation breeding.

Various technical support projects were maintained and measured. These include clonal maturation, clonal competition, pollen viability, and deer-resistance studies. Provenance trials were also maintained. Breeding continued for a deer-resistance heritability study.

4.6 Coastal White Pine

John King, Rich Hunt (CFS), Dave Pomeroy

In the fall of 2004 we established the first phase (with 80 families) of the F1 breeding program. Three sites were established – one on Texada Island and two on the Sunshine Coast. A second phase is currently growing at the Surrey nursery to be established in the fall of 2005. A third phase is to be sown in the spring of 2005. Parents for this effort came from the CFS screening program and were designated as either Slow Canker Growth (SCG) or Difficult to Infect (DI). SCG or "slow rusting" trees occur more frequently than DI. Also included as parents were some of the best of the Texada trees and some Idaho, Interior, and Dorena trees.

Although 2004 saw Rich Hunt's "official" retirement, luckily we have managed to retain Rich, part time at least, through an appointment within the CFS. Whether we will still have the services of Gary Roke waits to be seen. Ongoing pathology support by the CFS is seen as vital, especially with the establishment of the F1 generation. The university replacement that we see as

viable with the entomology support (see Sitka spruce) is not such a clear option with the blister-rust-resistance program.

We are still working with Stefan Zeglen on the regional deployment guide. This has been temporarily stalled because we hope to make it compatible with a similar guideline for Sitka spruce, but we hope to see it out early in 2006.

Recent publications on the white pine program include:

Hunt, R.S. 2006a. Blister-rust-resistant western white pine for British Columbia. Natural Resources Canada, Canadian Forestry Service Information Report BC-X-347.

Hunt, R.S. 2006b. Environmental and inoculum source effects on resistance of Idaho 72 western white pine in British Columbia. Can. J. Plant Pathol. 28:351-357.

Hunt, R.S., G.D. Jensen, and A. K. Dzonmbidoulah. 2006. Confirmation of dominant gene resistance (*Cr2*) in the U.S. white pine selections to white pine blister rust growing in British Columbia. Pp: 227-229 in Richard A. Sniezko, Saffiya Samman, Scott E. Schlarbaum, and Howard B. Kriebel, eds. Breeding and genetic resources of five-needle pines: growth, adaptability, and pest resistance: 2001 July 23-27; Medford, OR, USA. IUFRO Working Party 2.02.15. Proceedings RMRS-P-32. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 259 p.

King J.N. and R. Hunt. 2006. Five-needle pines in British Columbia, Canada: Past present and future. Pp: 12-19 in Richard A. Sniezko, Saffiya Samman, Scott E. Schlarbaum, and Howard B. Kriebel, eds. Breeding and genetic resources of five-needle pines: growth, adaptability, and pest resistance: 2001 July 23-27; Medford, OR, USA. IUFRO Working Party 2.02.15. Proceedings RMRS-P-32. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 259 p.

4.7 Interior Douglas-fir

Barry Jaquish, Jaz Ashley, Charla Phillips, Sonnie Hodge

Breeding: One hundred and thirty-five controlled crosses were completed, and 222 pollen lots were collected, processed, and stored for future crossing. Most of these crosses and pollen lots were from the high priority Nelson SPU.

Progeny Testing: Four 15-year-old sites in the Mt. Robson test series (Quesnel Lake SPU) were maintained and measured. Parental breeding values for individual tree volume (in percent gain over the average wild-stand population) at age 40 based on the measurement trait height at age 15 ranged from -62.6 to 41.8.

Armillaria Resistance Screening: The joint BC Ministry of Forests/Forestry Canada pilot project to screen interior Douglas-fir families for resistance to *Armillaria* root disease continues to produce promising results. To date, overall mortality attributed to *Armillaria* is 75 percent; however, there appears to be considerable variation in mortality among families (range 19-100 percent). Interestingly, there appears to be a positive relationship between mortality and elevation of the parent tree. This experiment will be terminated in fall 2005.

Site	Elevation (m)	Num. of Trees	Mean height (cm)	Survival (%)	Damage (%)	Range family of mean heights (cm)	Heritability (individual)	Heritability (family)
Kinbasket Lk	760	5797	428.7	87.6	2.6	356 - 571	20+/- 05	58+/- 15
FCTR	760	4798	462.4	72.6	8.5	286 - 571	30+/- 08	68+/- 12
Tete Jaune	914	2582	363.4	75.3	1.6	171 - 457	56+/- 08	77+/- 10
Valemount	914	4251	343.9	81.3	1.3	203 - 457	35+/- 07	62+/- 10

Figure 7. Summary of the 15-year results from the Mt. Robson interior Douglas-fir progeny tests (209 families) measured in fall 2004

4.3 Interior Spruce

Barry Jaquish, Val Kirkby, Charlie Phillips, Bonnie Ponsge

The interior spruce tree improvement program is the oldest program in the BC interior. It began in the mid-1960s with parent-tree selection and open-pollinated progeny testing in three seed-planning units. It expanded rapidly in the late 1970s to encompass other important seed planning zones in the interior. The program focuses on improving traits related to tree size, sweep, wood traits, and resistance to terminal weevil (*Pissodes strobi*). Through recurrent selection for general combining ability, it has progressed to the point where improved first-generation seed orchards are in full production for most zones, and second-generation full-sib progeny tests are in place for three seed planning units (SPUs). The emphasis of much of the recent breeding work is to maintain and measure the vast array of progeny tests, to bring together populations from the two programs to accommodate new and larger seed planning units, and to predict population responses to predicted global climate change.

Progress During the Last Fiscal Year

Breeding: Twenty controlled crosses were completed in the newly expanded Prince George SPU. This crossing program completes the amalgamation of breeding populations from seven smaller seed zones. Seeds for field testing will be sown in spring 2009. In addition, 13 crosses were completed and 25 pollen lots were collected from parents in the Nelson mid-elevation SPU. Twenty-one crosses were completed by Riverside Forest Products staff in the Armstrong Thompson–Okanagan seed orchard.

Progeny Testing: Four six-year-old and three ten-year-old progeny tests were maintained and measured in the Fort Nelson and Nelson SPUs, respectively (Figure 6). Ten-year measurements were recorded from the Thompson–Okanagan high-elevation progeny test at the Skimikin Seed Orchard. This test was subsequently terminated and removed to make way for establishment of the Fort Nelson spruce seed orchard. Wood relative density was determined by the oven-dry weight/green volume method on wood disks collected from 10 trees per family in the Okanagan–Adams progeny test at the Skimikin Seed Orchard (Figure 6). In the second-generation test series, the four six-year-old bulking/rating tests were maintained and measured, while the East Kootenay and Prince George sites were numbered.

In the support research program, three 15-year-old East Kootenay realized gain tests were maintained and measured, and five candidacy tests of *Lonatis* seedlings were maintained and measured. In the two-year-old wheat bed Prince George genecology study at Vanderhoof, shoot elongation was measured 21 times throughout the growing season. The three accompanying field sites near Prince George were also maintained. Seedlings for the new climate change study were also grown. Offsite and cold-stored and will be planted this spring on 18 climatically contrasting sites in BC, Alberta, and Yukon Territory.



Ov relative density at 19yrs

Figure 6. Relationship between growth trait parental breeding values for wood relative density (age 19) and individual tree volume at age 46 in Okanagan–Adams interior spruce at the Skimikin Seed Orchard.

Figure 4—Summary of results from the Pine Mistletoe and Mistletoe Interior Spruce programs, from inception to fall 1999.

Pete Hessburg presents Douglas-fir dwarf mistletoe results at the Western Forest Insect Work Conference to participants.

A Genetic Resistance for Western Gall Rust (Endocronartium harknessii) in Lodgepole Pine

Richard Sniezko, Jerry Kegley, Vicky Berger

Western gall rust is a serious disease pest of lodgepole and ponderosa pines throughout much of the range of both species. In lodgepole pine the disease gains entry to branches or main stems thru needles of the current year's shoot extensions. After infection, a gall forms on the branch or stem. If a disease gall grows more rapidly than the branch or stem, it will girdle and kill the appendage at that point. Although stem galls may not kill a tree, they often result in breakage of the stem at the gall, rendering the tree non-merchantable.

In our first cycle (generation) of lodgepole pine progeny testing across several SRIs our focus has been on improving stem volume growth. When measuring a particular progeny site, we have assessed damaging agents (diseases and insects) only if 10% or more of the trees were affected. Most of our 70 test populations are healthy and exhibit few disease or insect pest problems. Three of our thirty-nine first cycle test sites have shown western gall rust infection rates exceeding 10% of trees tested. Disease severity surveys of two progeny test sites north of Mackenzie in 1994 (with the technical guidance of Richard Hirsch, northern region pathologist) are providing insights into patterns of genetic resistance to western gall rust in lodgepole pine.

Although most of the progeny tested parents are from natural stand origins, our tested parents from a range of 10 to 60 degree north, 120 to 150 degree west, and from a wide range of elevations. Disease resistance as measured by the mean number of galls per tree varied widely with geographic variation of latitude, longitude, and elevation of parent tree origins. Although some provenances exhibited greater average levels of resistance than others, the variation in resistance among families within provenances was the major source of variation in disease resistance (see Figure 45). Family resistance values measured on tree vigor were highly positively correlated, suggesting little trait cost (see Figure 11). Apparent high heritability for gall rust resistance and low cost suggest the possibility of selecting for a rust-resistant seed orchard for this high rust hazard area in the north central interior. There appears to be no correlation between rust resistance and breeding value for growth (see Figure 45). Selecting solely for rust resistance (i.e., no improvement in growth) may be the best strategy for the Mackenzie area.







4.12 Western Larch

Barry Jaquish, Val Ashley, Cicero Phillips, Bonnie Ralough

The BC western larch program was initiated in 1957 and has advanced rapidly. The program strives to improve tree size through recurrent selection for general combining ability.

The main components of the program are phenotypic selection of parent trees in wild stands, open-pollinated progeny testing, roll-based seed orchards, and second-generation crossing. To date, 930 parent trees have been selected and wild-pollinated families of 827 of these parents have been established in progeny tests across 34 sites in two SPUs (East Kootenay and Nelson).

Two grafted seed orchards were established in 1980 and are now in full production. Presently, over 70% of the 4.3 million western larch seedlings planted annually in BC originate from Class A seed orchard seedlots. Second-generation crossing is over 60 % complete.

Breeding: In the second-generation crossing program, 30 and 34 crosses were completed in the East Kootenay and West Kootenay SPUs breeding orchards respectively. Eighty-eight pollen lots were collected, processed, and stored for future crossing.

Progeny Testings: Four 10-year-old sites in the East Kootenay Series II test were maintained and measured. Parental breeding values for individual tree volume (in percent gain over wild-stand seedlots) at age 40 based on the measurement trait height at age 10 ranged from -35.8 to 22.8.

Realized Gain Testings: Three realized gain test sites were established in the Nelson SPU. The main objective of these tests is to estimate yield of western larch seedlots of different genetic quality on an area basis. The experiment was established in a randomized complete-block design with three levels of site quality (low, medium and high), four levels of genetic quality (composite elite full-sib crosses, high genetic-worth seed orchard, low genetic-worth seed orchard, and composite operational controls), three levels of spacement (1.5 x 1.5 m; 2.5 x 2.5 m; and 3.5 x 3.5 m), and two replicate blocks per site. Plots consisted of 144 trees planted in 12 x 12 tree configuration.

Site	Elevation (m)	Num. of trees	Mean height (cm)	Survival (%)	Damage (%)	Range family of mean heights (cm)	Heritability (individual)	Heritability (family)
Wigwam Cp	1140	4149	673 X	85.7	5.7	530 - 923	.27+/- .06	.77+/- .10
Lower Lardeau Cp	1140	4577	468 Y	76.1	1.3	258 - 678	.35+/- .07	.69+/- .10
Gerdie Cp 4	1470	4999	674 X	83.8	2.8	258 - 952	.61+/- .08	.93+/- .10
Upper Lardeau Cp	1270	5189	561 Y	88.1	1.9	527 - 762	.25+/- .06	.58+/- .10

Figure 19. Summary of the 10-year results from the East Kootenay Series II western larch open-pollinated progeny tests (*Will Hamilton, measured in fall 2000*)

5.0 Operational Tree Improvement – The Eighth Year in Review

Roger Painter

In many ways this has been a watershed year for this program. Over the years, the Operational Tree Improvement Sub-Program (OTIP) has provided the focus for investing in our orchards for the purposes of increasing genetic gain as well as seed yields. Seed requests in 2004-5 have now reached 50% of provincial reforestation needs, and our genetic gain is close to meeting the overall target set for 2007 by the Forest Genetics Council in 1997. This sub-program represents the strategic device that we have employed to invest in existing orchards and bring them up to their potential. This, coupled with the development of replacement stock from the breeding programs, has raised the genetic worth of orchards to more acceptable levels. At the same time, this program relies on a system of performance management to monitor progress and set reasonable targets for project success. Orchardists and researchers have responded to these approaches and in many instances have over achieved.

Although 2004-5 was a lighter than normal year for most species crops, considerable work was done in stock replacement and studies on seed supply issues. Unfortunately, some crops sustained large attacks by insects, specifically *Dioryctria*, that reduced yields. This underlined the need to focus more attention on Seed Pest Management, along with establishing a stronger mandate for research in that field and further development of the Seed Pest Management Technical Advisory Committee and its associated sub-program. Future research and development projects that used to come through OTIP will now be sent to that sub-program. This is an example of the continued definition of focus and planning procedures within the overall program that has allowed us to concentrate our efforts despite limitations in funding. The Forest Investment Account continues to show strong support for our program and to keep funding constant at current levels. Available supplies of seed from our seed orchards continue to increase, and the genetic quality of seed crops continues to improve. In the long term, our province will enjoy a future supply of excellent wood products and a healthy resource as a result of the foresight of investing now.

In 2004-5, the two technical review committees reviewed a total of 76 requests for funding in OTIP. The total amount of proposals received was approximately $930,000. The Coastal and Interior technical committees recommended that $694,813 be approved. The breakdown of investment by region for all of OTIP is as follows:

Interior projects	44	$456,837
Coastal projects	30	$238,797
Overall total	**74**	**$694,813**

Figure 14. Number of projects and funding by region

The Forest Genetics Council has the support of various species committees that review annual projects and develop priorities. These reviews give direction to both the Tree Breeding and the Operational Tree Improvement programs. Similarly, the Extension and Gene Conservation TAC's have developed their own priority review procedures. Tree Breeding, as a Ministry of Forests responsibility, is not part of the OTIP "Call for Proposals" process. Priorities for the Tree Breeding sub-program are still developed through the species sub-committees, but decisions on projects to be undertaken are reached through co-ordination with the Tree Improvement and Research branches of the Ministry of Forests and the FGC Program Manager. Considerable investment is needed in this sub-program, which will continue to receive strong levels of support to ensure the development of new production stock for future orchards. New orchards will focus on producing seed for Seed Planning Zones in which production capacity is low and/or priorities for genetic- quality seed are high. These new orchards will also help produce stock to replace older, less advanced stock in existing orchards. Work in Tree Breeding has already provided much of the genetic material to establish a number of new orchards recently and to enhance existing ones. The development of a long-term investment program through SelectSeed Ltd. has produced over 35,000 grafts for new orchard establishment so far.

Technical support within the OTIP sub-program is an integral part of tree improvement in general and provides an excellent avenue for operational problem-solving. The Lodgepole Pine seed-set issue has been an excellent example of focused investigations to solve serious problems affecting tree improvement. Listed below is a project breakdown by areas of investment for 2004-5:

Tree Breeding	36
Gene Conservation	7
Operational Production	57
Technical Support	17
Seed Pest Management	4
Extension and Communication	4
Seed Planning and Information Tools	2

Figure 15. Number of projects by areas of investment

The tree improvement industry represents a broad spectrum of partners, including forest companies, the provincial government, the Canadian Forest Service, universities, private bio-technical companies, and individuals. In 2004, the Tree Improvement Program involved 37 separate contributors from all parts of our industry. The Forest Genetics Council continues to provide sound leadership and management through its technical committees and through the use of a strong business management approach for setting priorities and goals for investment. It also offers a strong sense of renewal through review of program goals and strategic planning.

5.1 Orchard Projects

5.1.1 Saanich Forestry Centre

WFP

Annette Van Niejenhuis

Western Forest Products Inc. manages production tree seed orchards for eight seed planning units, together with hedge orchards for one seed planning unit at the Saanich Forestry Centre. As a co-operator in the Forest Genetics Council programs aimed at delivering quality seed in quantity to the coastal forest regeneration programs, WFP has employed OTIP funds for incremental management techniques in six seed planning units.

SPU 0107:

Our low-elevation coastal Douglas-fir orchard is being improved with replacement clones. One hundred and seven ramets of fourteen clones have been acquired (ten new clones and four orchard clones). These ramets will be planted in the orchard, replacing mortality and very poor genetic quality clones. Aggressive roguing and pursuit of gain is deferred until more seed is banked in this SPU. With the addition of these ramets, the average breeding value of this orchard is raised from 11.1 to 12.1.

Only a small crop was available for management in 2004, and insect damage of cones was the highest ever recorded in this orchard. For these reasons, crop management was minimized. High-gain pollen (6.2 litres) was harvested, extracted, and banked for future deployment.

SPU0206:

An improved western redcedar replacement orchard was planted in 2004. With this stock, the average breeding value is raised from 5.4 to 6.5. This orchard includes clones from the first three series of western redcedar polycross progeny tests. As additional results from these trials are received, about 50% of the ramets will be rogued and replaced with clones from the final three series of progeny tests, with the expectation of improving the breeding value to 10 or higher.

The 2004 western redcedar crop was infested with cone midge. Trees with significant crop were treated with dimethoate under SUP 111-210-04/04. On sprayed trees all midge were killed, but few seeds developed and no crop was harvested. It is unknown whether the treatment, the damage from the midge prior to the treatment, or

other factors such as poor pollen success caused the poor seed set. The treated trees showed foliar damage.

SPU0304:
Supplemental mass pollination was applied to the western hemlock crop, focussing on clones with a high propensity to self-pollinate. The resulting seedlot has a genetic worth of 14 and is estimated to produce 110 thousand seedlings. High-gain seed is needed in this SPU; little seed of genetic worth of 9 or greater is banked.

SPU0601:
Seed production in the weevil resistant Sitka spruce orchard continued to increase. Supplemental mass pollination to improve seed set was applied. The resulting crop, including surplus seed from past controlled pollination, has a weevil resistance rating of 86% and is being deployed to the 2005 nursery crop for 74 thousand trees.

Ramet replacement will continue with the addition of 31 ramets. Our intent is to include all clones for which testing is underway. When test results are known, roguing will maintain high weevil resistance, combined with volume selection.

SPU1104:
More than 900 donor stock replacements were established in the yellow cypress production hedges. With these replacements, the average clonal value of the hedge is increased from 19% to 20%. Spot treatments for trisetacus mites were required to maintain cuttings production, and pruning of young hedges was implemented to improve the quantity and quality of the material. The 2005 nursery crop includes 314 k of yellow cypress cuttings with a clonal value of 19 or greater.

SPU1110:
A study of the optimum rooting strategy for technical support in deploying yellow cypress was initiated. Cuttings from the TimberWest and Western Forest Products hedges harvested and set at Sylvan Vale and Arbutus Grove nurseries on four dates will provide information about optimum timing for setting. The expectation is that optimum timing will result in improved nursery recoveries. This will lead to maintaining present stock prices and prevent increased stock prices. It will also improve the deployment of the high-gain material available through this clonal regeneration program.

SPU1112:
A clonal rootability study of the MoF high-gain yellow cypress clones will permit selection of good rooters in the production hedges. A selection of nine MOF clones from TimberWest and Western Forest Products hedges and twenty-two WFP clones from Western Forest Products hedges were set clonally in November of 2004 at Arbutus Grove and Sylvan Vale nurseries. These trials will be repeated to determine which high-gain clones are poor rooters (<50% success). These will be removed from production hedges to improve the overall nursery recoveries in the yellow cypress rooted-cuttings program. Those poor rooters with extremely high clonal values will be included in breeding orchards to determine if rootability is heritable and if breeding can alleviate the poor rootability and lead to new high-gain selections with high rootability.

SPU2401:
No crop was available for management in the high-elevation western hemlock orchard. Ramets were acquired for replacement and are maintained, but the planting is deferred until site problems are addressed.

Western Forest Products has been tracking the genetic worth of SelectSeed deployed to the nursery (Figure 16). Significant gains in the average genetic worth of yellow cypress have been achieved since 1996, with the 2004 nursery crop averaging 20% volume gain. Douglas-fir and western hemlock also display strong gain trends in this period. Western redcedar orchards will soon be delivering significant volume gain to the regeneration programs. The Sitka spruce program is geared to weevil resistance ahead of volume; thus, it will be a decade before volume gains are delivered there. The weighted average volume gain of all A-class SelectSeed deployed by Western Forest Products is now above 6%. Factoring in the wild stand seed deployment with the A-class SelectSeed deployment, average volume gain drops to just under 5%. Throughout this period the deployment level has increased from 65% to 75% across all of our coastal regeneration programs, with 100% deployment in most seed planning units for which orchards exist. The Forest Genetics Council's Operational Tree Improvement Program assists the development and production of high-quality seed for deployment in our coastal regeneration programs.



Figure 16. Weighted Average gain for SelectSeed in WFP orchards only



Plate 6. Yellow cypress clonal trial selections

Plate 7. J. Russell discussing wood quality and breeding at Coastal Silviculture Committee meeting

Plate 8. Our first weevil resistant seedlot from Ss 172

5.1.2 Sechelt Seed Orchard

CANFOR

Patti Brown

Douglas-fir

Orchard #177 produced its first crop in response to the 2003 inductions. Although 10 litres of cones were collected from controlled cross-pollinations, only 1 kg of seed with a GW of 18 was extracted. Despite being bagged during pollination and after for insect protection, *Dioryctria* must have been present on the branches prior to bagging, and most of the crop was destroyed.

One hundred fifty ramets in orchard #116 were induced in June 2004 in response to a lack of availability of high-elevation Douglas-fir seed.

Western redcedar

The ramets in the holding beds were fertilized monthly and pruned to increase the surface area. Those parents with a GW of >6 were induced, along with the main orchard (#186), in July. Orchard #186 and the holding beds were rogued in February to remove ramets from Series 2 with age 7 results of GW<5. Pollen from the top ten parents was collected for reapplication to females in orchard #186 and seed-producing ramets in the holding beds. The sunny dry weather in February 2005 caused the males to shed pollen early and the females to lag behind, resulting in less selfing but also less seed set from natural pollinations.

Project SPU0211 advanced the breeding program material into orchards by grafting Series 3 and Series 4 material. In November 2004, 5,292 grafts were completed, and these will be incorporated into orchard holding beds in 2005.

Western white pine

The goal of SPU0801 was to increase the slow-canker-growth resistance value of orchard #174 in the long term and to produce a crop with > 50% major gene resistance (MGR) in the short term.

This year's MGR crop (> 50% resistance) was monitored and closely spot treated for *Leptoglossus* from May to August. The resultant high-yield crop has the potential to produce 300,000 plantables. The 2005 crop was hand pollinated with either MGR pollen or an elite mix of slow-canker pollens.

Eighty lower-resistance-value trees were removed from orchard #174 in the summer of 2004, and they will be replaced with the survivors of the 245 higher-value, slow-resistance clones that were grafted in April 2004.

Yellow-cedar

SPU1101 continues with the goal of producing 300,000 stecklings with a GW of > 12% annually for our coastal operations. To achieve this, donors are grown and cultured at Cairnpark Nursery. The top selections from the Ministry of Forests Series 2 and 3 breeding programs plus Western Forest Products' breeding program were obtained in 2004/05 and are being cultivated at Cairnpark to produce large numbers of future donor plants, thereby increasing the average GW value of our vegetative lot from 12 to 18.

5.1.3 Mount Newton Seed Orchard

TimberWest

Tim Crowder

Seven Operational Tree Improvement projects (OTIP) were carried out at Mount Newton Seed Orchard this year.

Enhancing Productivity and Gain in Four Coastal Douglas-fir Orchards (SPU 0106)

SMP was carried out in orchard #183, and on early- and late-flowering clones in orchards #134 and #154.

Cones were monitored for *Contarinia* midge during the pollination season, and because infestation levels were high, all three orchards were treated with dimethoate.

Over 800 ramets were induced by stem injection of GA4/7 to produce a crop in 2005. This included some ramets from orchard #197 for the first time.

Following the harvest of the 2004 crop, 40 ramets with low BV were rogued from the orchards, and 250 grafts were planted in vacant orchard locations. Grafts that were too small to plant out were maintained in a holding bed.

In the fourth quarter, 1,200 grafts of high BV clones were made for future upgrading of the Douglas-fir orchards.

Enhancing Productivity and Genetic Gain in Two Western Redcedar Orchards (SPU 0205)

The genetic worth of the 2004 seed crop in orchard #140 was 7%, the highest gain so far for an open-pollinated seedlot from this orchard.

Upgrading programs continued in 2004 in orchards #140 and #152. Vacant spots in both orchards were de-stumped and planted with high-ranking clones from the holding beds. Top-ranked clones from all three test series were included.

Orchard #152 was induced for a seed crop in 2005 through foliar application of GA3.

Increased Production and Gain from Western Hemlock Orchard #182 (SPU 0310)

Two clones with BV less than 4 were rogued from orchard #182 and replaced with a total of 30 ramets from several clones with BV of 15 or greater. An additional 100 ramets were grafted to provide replacement stock for future roguing operations.

Because the ramets in orchard #182 are still relatively small, SMP was used to supplement the insufficient pollen cloud. The genetic worth of the 2004 seed crop in orchard #182 was 14%.

One-quarter of the trees were large enough to be induced for the first time, and so were injected with GA4/7 to produce a crop for 2005.

Management of Western White Pine Orchard #403 (SPU 0808)

Three hundred ramets were transplanted from the holding bed into empty spots in orchard #403, and 300 grafts from the previous year were transferred into the holding bed. Two hundred new grafts were made of clones that had failed to take or were not available previously.

MGR pollen was obtained from CanFor and applied to most of the developing cones for the 2005 seed crop.

Leptoglossus bug continued to be a threat in 2004 and required frequent monitoring and control.

Maintenance of *Abies amobilis* Orchard #129 (SPU 0901)

Because orchard #129 is the only remaining Ba orchard in the province, the objective of this project is to maintain and manage the orchard to increase the knowledge base for seed production in this species. In addition, Ba seed for reforestation is produced.

2004 was a light crop year for orchard #129, following the exceptional crop of 2003. The 1,115 ramets were maintained with irrigation, pruning, and vegetation control.

Yellow-cedar Hedge Management (SPU 1109)

The objectives of this project are twofold: 1) to continue to produce 300,000 cuttings annually with a GW of 12% or greater, and 2) to bulk-up and hold 120 clones from series one and two of the MoF testing program.

These objectives were achieved through maintenance of the productive field-based hedge of 3,000 donor plants, and maintenance of 2,000 potted ramets in a greenhouse.

Management of High-elevation Western Hemlock Orchard #130 (SPU 2402)

A crop for 2005 was induced by stem-injecting non-crop trees with GA4/7. The 2004 crop had a GW of 6%. Cones on non-tested trees in the orchard were not harvested.

5.1.4 Bowser Seed Orchards

Dan Rudolph

Genetic Upgrading and Enhancing Seed Yields and Quality From Second-generation Douglas-fir Seed Orchards (SPU 0110)

This project is a multi-year effort and is being conducted in three second-generation seed orchards, #149, #162 and #168 (a micro-orchard containing the same clones as #162), at the Ministry of Forests' site at Bowser.

There are two main objectives:

1. To upgrade the genetic composition of the orchards by roguing the lowest-ranked clones and replacing them with higher-ranked parents, and
2. To enhance the seed yield and genetic worth of the seedlots produced by:
   -using cone-induction techniques on all suitable candidates in each orchard.
   -using controlled-pollination techniques in both orchards in order to produce elite seedlots.
   -using orchard-management techniques to optimize growing stock vigour and crop health and so maximize yields.

Results

1. Upgrading

At the Cowichan Lake Research Centre, 2,451 grafts from 17 of the highest-ranked clones (breeding values between 16 and 26) have been propagated. Twelve hundred of these ramets will be transplanted into a holding area on-site during the fall of 2005; the balance of the stock is to be transplanted into a similar holding bed at the Saanich site for eventual inclusion in an orchard to replace orchard #149, which is currently at Bowser. The holding area at Bowser is to be developed during the fourth quarter of this fiscal year.

2. Enhancing seed yield and genetic worth

Controlled crossing: 89 ramets from 17 of the highest-ranked parents in each orchard were identified, and isolation bags were put on 1,030 branches of these ramets. Controlled crosses using pollen mixes with high genetic worth were made. Pollination bags were replaced with insect bags after orchard receptivity had passed.
Induction: Suitable candidates for induction were identified in each orchard. Criteria used to establish a candidate list included negligible current year's crop, no crop from last year, no induction treatment last year, sufficient vigour, and an adequate number of cone-bearing sites. Induction treatments consisting of a double overlapping girdle and application of fertilizer at the rate of 300 kg N/Ha were applied. Irrigation was withheld until mid-July in order to increase drought stress on the trees.
Orchard management: All orchard trees were maintained through appropriate cultural practices.
Insect management and control: surveys were conducted for *Contarinia oregonensis and Leptoglossus occidentalis.*

Output and deliverables

1. Upgrading

The removal of the lower-ranked clones in orchard #162 during the first year of this project raised the overall genetic worth of the orchard from 12.5% to 15%. When the new clones are established in the orchard, the overall genetic worth of the orchard will rise to 20%.

2. Enhancing seed yield and genetic worth

Controlled crossing: Seedlot #90305 was produced from the controlled crosses done in spring 2004. Seed yield was 0.464 kgs of seed with a genetic worth of 18.6.
Induction: A total of 275 trees were induced – 62 from orchard #149, 138 from orchard #162, and 75 from orchard #168. Bud surveys to rate efficacy of the treatments are to be done in late March 2005.
Orchard management: Foliar nutrient samples were taken; fertilizer was applied for both growing stock and crop maintenance; irrigation was applied when necessary; trees were basal-pruned, top-pruned, and inter-nodal-pruned, and identity tags were upgraded prior to cone collection.
Insect monitoring/control: Because *Contarinia* levels were within the acceptable threshold level, no control measures were necessary. *Leptoglossus* levels were also low; therefore no control measures were undertaken.

5.1.5 Kalamalka Seed Orchards

Chris Walsh

Ten OTIP projects were approved under the operational production sub-program for 2004/2005 at Kalamalka Seed Orchards. The funding allowed for a significant enhancement of the effectiveness of our orchards in delivering improved seed. Activities included:

- Improving orchard composition through grafting higher-breeding-value ramets, maintaining recently grafted high-value ramets destined for orchards, planting rootstock for future grafting, transplanting the older higher-value ramets to the orchards, and roguing lower-value ramets from the orchards;
- Improving orchard seed quantity and quality through pollen management, including collection of high-breeding-value pollen from clone banks and the application of Supplemental Mass Pollination; and

- improving orchard productivity through pest management and other management activities.

Pest management activities funded included:

- use of Safer's Soap sprays to control pales weevils,
- removal of weevil-infested spruce leaders to reduce weevil populations,
- removal of pine pitch moths damaging orchard tree stems,
- baiting for control of rodents feeding on tree roots,
- sanitation picking of cones in orchards with non-collectible crops to reduce pest populations,
- dormant oil application to control larch adelgids,
- sprays to control *Cydia* and *Strobilomyia* in Sx cones,
- sprays to control *Leptoglossus*, and
- sprays to control spittle bugs and *Zeiraphera* in lodgepole pine orchards.
- other funded management activities to boost productivity and gain included foliar analysis to determine the nutrient status of orchard trees and crown management of orchard trees.

The OTIP funding was instrumental in increasing both the quantity and quality of seed produced. At Kalamalka in 2004 we produced approximately 180 kg of interior spruce, western larch, lodgepole pine, and western white pine seed, equivalent to over 25 million seedlings, with an average GW of +16. Kalamalka seed is being used over large areas of the interior of the province.

Project	Species	SPZ	Orchard	Pollen Collected (litres, dry)	Trees Pollinated
SPU0401	Sx	NE	305	3.0	961
SPU0502	Sx	NE	306	3.0	496
SPU0701	Pl	NE	307	3.0	1,987
SPU1302	Lw	NE	332	1.0	678
SPU1501	Pw	KO	335		1,455
SPU2201	Fd	NE	324	1.0	
SPU2601	Sx	IR	405	2.0	321
Totals				16.0	5,261

Figure 17. Pollen management activities by project

Project	Species	SPZ	Orchard	Rogueing	Grafts Made	Maintained	Roguebank	Transplants
SPU0401	Sx	NE	305	61	150	336	120	210
SPU0502	Sx	NE	306	40	200	277	150	128
SPU0701	Pl	NE	307			46		
SPU1302	Lw	NE	332	30	150	402	150	171
SPU1501	Pw	KO	335		28	236	150	41
SPU2201	Fd	NE	324			87	120	77
SPU2501	Sx	EH	304	158				20
SPU2201	Pl	EH	340		126	276	120	16
SPU2401	Lw	EH	333	266	150	367	120	136
SPU2601	Sx	IR	405		80	247	90	32
Totals				574	913	2183	970	823

Figure 18. Orchard composition activities by project

5.1.4 Vernon Seed Orchard Company (VSOC)

Tim Lee

The seven funded projects approved for this year were an important part of our program, funding enables our staff to undertake projects that promote the complete management of all one and a half generation production orchards on site. Young orchards require additional activities to keep orchard seed production at its highest level each year. Our mature orchards that required upgrading were funded for the roguing of lower breeding value or production challenged families, and now the grafting and planting of replacements for the benefit of the Seed Planning Unit are completed.

VSOC manages eight orchards for production. These orchards were the first phase of our site development. The second phase included the addition of five more orchards. These orchards are not expected to be a part of the seed supply for eight to ten more years.

Individual Items

- **Pollen work** – Our young orchards have a limited supply of natural pollen, and with the processed frozen pollen from previous years, we can aid the early or late flowers. VSOC historically has picked, applied, stored, and frozen many litres of pollen for all our orchards. Our mature spruce orchards, on the other hand, are now left to the natural pollen cloud that has developed.
- **Supplemental Mass Pollination (SMP) treatments** – SMP is used in our orchards. The timely application of pollen can make the difference in ensuring the production of seed. Helicopters have proved to be a key factor in pollen management in large young orchards where pollen clouds are only beginning to be detected. Our pollen applications are getting more pollen into the receptive female flowers. Previously we were receiving only marginal pollen uptake into pollen tubes of pine.
- **Grafting** – Timely seed replacement is dependent on an active, well-planned program that is aware of the needs of each SPU. Rootstock holding beds are managed with regard to a projected need for the continued improvement of each orchard. Information on family rankings and the advancement of the next generation of selections are always in the back of each orchard management team's mind. Orchards evolve as the breeder and orchard management teams see the next generation of seeds included into the orchard, but the team is always mindful of how to manage the transition periods that affect seed supply from each orchard.
- **Rogue replacement** – Replacement of rogued positions within an orchard are also important to orchard development.
- **Insect and disease control** – An important part of seed production is the control of such insects as cooleyinous scoliidentalis, European pine shoot moth, pine pitch moth, and adelgids. These are a few of the insects that are monitored and controlled for the protection of our orchards' health and production.

Activities undertaken each year affect the quantity and quality of seed produced for this province. Funding made available has had a positive effect on the Forest Genetics Council's goals of 12% gain and 75% use of Class A seed for our province.

Break down of projects

Vernon Seed Orchard Company (FCP 2000)

SPU Project#	Species	Orchard	Pollen Units	GMP - Treatments	Insect & Disease	Cone Made	Grafting Scion	Planted
1234	AB	Site	Business Analysis for all Production Orchards					
1345	PL	223	7	2000	2488	1200	1300	1400
1354	PL	218	25	5000	5800	600	600	667
1901	PL	218	12	3000	4000	200	250	275
2002	Sw	226	8	200	221			
4123	Sw	225	4	200	180			
4501	Sw	241	15	800	1020			
Totals			82	11400	14409	2000	2300	2742

Figure 16: Break down of projects

[redacted] Sagaluna Seed Orchards

[redacted]

Sagaluna seed orchards are located on Sagaluna Beach, about 60 km southwest of Mintarong. At this site, three subgaponi pine orchards and one Douglas-fir orchard received STIP funding for activities to increase the yield and genetic gain of seed produced. Orchards 008 and 011 provide seed for the Mt Thompson-Sibanagan low-elevation seed planning unit (SPU), orchard 012 provides seed for Mt Mahon low-elevation SPU, and orchard 021 provides seed for Mt Mahon low-elevation SPU.

All orchard 008 is a provenance-based orchard with an estimated genetic gain of 0%. All orchards 011 and 012 are 1.5 generation orchards each with an estimated genetic gain of about 10%. All orchard 021 is a 1.5 generation orchard with an estimated genetic gain of 20%.

In 2006, STIP funded a number of activities in these orchards. STIP projects 0705, 0801, 1002, and 1101 covered activities in orchards 011, 012, 008, and 021 respectively. These activities included insect and disease control, rodent control, cover management, foliar analysis, pollen monitoring, pollen collection, and supplemental mass pollination (SMP).

[redacted] Orchard 008 activities

In early spring 2006 foliar samples were taken to determine the appropriate fertilizer mix for spring application. Foliar samples were taken again later in the season for fall fertilization. These analyses helped direct fertilizer applications to ensure optimal nutrition.

Pollen monitoring began in May with monitors set up within all three Mt orchards and between orchards. Pollen monitoring indicated that there was a substantial pollen supply in orchard 008, and therefore SMP was not conducted in this orchard. The natural pollen cloud was much smaller in orchards 011 and 012, but SMP was not conducted due to operational limitations.

No protandry was observed in any of the Mt orchards; the shedding of pollen and flower receptivity coincided well. As the volume of pollen being shed increased, an air-blast sprayer blowing a very fine water mist was used on calm days to ensure good distribution of pollen in the orchards.

During the pollen flight period, pollen was collected from orchards 011 and 012 for future use. Whole pollen buds were manually collected by cone cutting and brought into the operations for drying and processing. Over 10 litres of pollen with an average IMC of 60 to 70 was put into storage.

Throughout the season, all Mt orchards were monitored for insect, rodent, and disease problems. This ensured that measures were taken to protect seed health and seed quality. Herbicide sprays were applied as necessary to control cyperaceous weed crop. Sprays indicated that fungicide sprays were not required in 2006. Poison baits were set out to control rodents. Bashing on male buds, and pinch shifts were removed by hand.

A small amount of cone pruning in each orchard encouraged branching and maintained the trees at a manageable height without compromising flower production.

[redacted] Orchard 011 activities

Pollen monitoring began on the first flight of pollen in the orchard. Pollen monitors were erected in 11 locations within the orchards and they were monitored daily until the pollen flight was completed. Pollen cloud and female receptivity were monitored throughout the pollination period, culminating the timing for SMP applications.

Pollen was applied three times in April with stored pollen from 2005 collections (previous STIP projects). Pollen for future use was collected by 11 PSP crew at the Kalawaska Forestry Centre. Trucks in Mercy Airport and Inland Airbay for providing access to their cone banks for pollen collection and packing by PSP crew. The pollen was processed and dried at PSP and put into freezer storage. Approximately half (~3 litres of pollen) is now available for SMP efforts in 2009.

With the exception of SMP in the Mt orchards, all projects were completed as planned. In the Mt Thompson-Sibanagan (~ha, 10 ~7 kg of seed was collected in 2006, with the potential to produce ~11 million seedlings. In the Mt Mahon (~ha, 5.5 kg of seed was collected, with the potential to produce ~7.7 million seedlings. The Mt Mahon (~ha orchard produced a small crop of approximately 100 grams of seed with the potential to produce 0.060 seedlings.

[redacted]

[redacted]

[redacted]

a rust resistance screening trial, and a site was prepared for about 3,500 more to be planted in 2005. The Aiken garden was maintained and white pine seedlings were inoculated in September. About 2500 trees from a Douglas-fir genecology trial were thinned and mulched. A total of 9,130 trees were removed from plantations no longer required, and a site was prepared for the planting of a spruce provenance test in 2005. The plantations were the main portion of the Canadian Tree Improvement Association (CTIA) summer tour here in July.

Photo 16. Trees being removed from old plantations

Photo 17. Preparing the site for a spruce provenance trial

Photo 18. Spraying in the Bulkley Valley spruce orchards

Photo 19. Mike Carlson discusses the results of birch provenance trials at the Skeena Seed Orchard with CTIA participants

Photo 20. Planting replacement grafts in the holding bed

5.1.11 Saanich Seed Orchards

Carolyn Lohr

Four Operational Tree Improvement Projects (OTIP) were carried out at Saanich Seed Orchards this year.

Establishment and Monitoring of Regional Pollen Monitors in the Saanich Peninsula (SPU 0109)

Five, seven-day recorders were established at locations throughout the Saanich Peninsula. The objective of this project was to provide consistent estimates of local pollen contamination to Douglas-fir seed producers in the area, in order for them to determine the effect on the genetic worth of the seedlots produced. Pollen flight was monitored from March 30 to April 26. Data were counted and averaged, and counts were forwarded to local Douglas fir seed producers.

Genetic Upgrading and Enhancing Seed Yields and Quality from Rust-resistant White Pine Seed Orchard #175 (SPU 0804)

The objectives of this project were twofold: to re-graft and re-establish orchard #175 up to pre-vandalism numbers with genetically upgraded stock and to operationally manage the existing orchard trees to increase the yield of filled seed.

Fourteen new clones consisting of 238 ramets were grafted at Kalamalka, and 180 ramets from CLRS are now being maintained in holding beds at Saanich Seed Orchard. Larger ramets will be planted into empty spots in orchard #175 in spring 04 or fall 05.

To increase yields of filled seed, bud and phenological surveys were conducted. Three litres of pollen were collected and re-applied to female flowers. Collection of 15,187 cones (45.2 Hl) yielded 23.1 kg of seed (results of 2003/04 SMP). This is an average of 70 seeds/cone, which is considerably higher than the 50 seeds/cone if no SMP is applied.

Two hundred and seventy five trees were maintained by irrigating, fertilizing, and controlling insects and vegetation.

Orchard Management Operations to Maintain Productivity and Increase Gain in Orchard #181 SM FDC(SPU 1902)

Orchard #181 is a 95-clone 1,037-ramet, first-generation SM, mid-elevation Fdc orchard with progeny being planted in spring 2005 for testing. Five-year measurements will be available in 2009/2010. At that point in time, the orchard will be rogued to a BV greater than 5.

Management operations included surveys for health and vigour, graft maintenance, rootstock pruning, mortality replacement, fertilizer and irrigation application, foliar nutrient sampling, and pest surveys for foliage and cones. Pollen was collected and stored. The crop was not managed due to low numbers of female flowers and high numbers of insects.

Genetic Upgrading and Enhancing Seed Yields and Quality from High-elevation Western Hemlock Orchard #196 (SPU 2403)

Orchard #196 consists of 12 clones (79 ramets) of high BV moved from Campbell River Seed Orchard #143 and 12 clones (172 ramets) planted this season of high BV material from CLRS. The mean BV of this orchard is now 11. Ramet numbers were increased to reach effective Ne and high BV.

To increase seed yields, bud and phenological surveys were conducted. Three hundred and thirty-three mls of pollen were collected and re-applied. Three hundred and seventy-three grams of seed (62.5 litres of cones) were collected. The 2004 seedlot was blended with the 2003 seedlot because Ne was not high enough in 2004. The resulting seedlot has an Ne of 10 and a BV of 8%.

Management of the orchard included surveys for health and vigour, graft maintenance, rootstock pruning, mortality replacement, fertilizer and irrigation application, foliar nutrient sampling, and pest surveys for foliar and cones. A small N fertilizer/seaweed extract (kelp) application study was conducted for control of woolly adelgids over the 2004 season. Levels remained low, and because natural predators were present, no pesticide control was necessary.

5.2 Technical Support Projects

5.2.1 Cone and Seed Pest Management – Interior Operations

Robb Bennett

This project describes operational activities for the Interior seed pest management biologist, including all expenses associated with lab, field, and office work. Included in this description are activities for one co-op student (one term) and auxiliary technical support (as needed) to assist with operational pest management trials, data analysis, and development of new operational techniques.

Between April 1, 2004 and 31 March 2005 the Interior seed pest management biologist and the co-op student performed the following services.

1. Performed – 200 site visits, pest surveys and identification, and damage predictions and assessments.
2. Delivered – 50 written pest-survey reports to orchard managers and other seed production personnel.
3. Performed – 40 other pest identifications and management recommendations to Ministry of Forests personnel.
4. Dealt with – 25 requests for information from members of the public.

Work continued on in-house and collaborative development of protocols for seed bugs, seed wasps, coneworms, mountain pine beetle, and other insects of importance to Interior cone and seed production. In summary, the Interior pest management biologist:

1. initiated, continued, or completed seven in-house pest management research projects,
2. collaborated with university, government, and other research personnel on five other projects,
3. assisted in the preparation of mountain pine beetle risk assessment documents for PGTIS and Skimikin seed orchards, and
4. assisted in creating post-harvest management protocols for coneworm-infested cones.

As well as training provided for biology co-op students and orchard technicians, the Interior pest management biologist provided the following training experiences:

1. nine extension education presentations to grade school, college, and university students,
2. two professional presentations – Inland Empire Tree Improvement Cooperative, Canadian Tree Improvement Association, Superior Woods Tree Improvement Cooperative, the Forest Genetics Council Species Committee (two), and the Forest Genetics Council (two),
3. four pest management workshops for orchard staff, and
4. numerous "tail-gate"-type extension presentations to operational personnel.

The following publication resulted from this project.

Strong, W.B. 2004. Seed Orchard Pest Management and Insecticide Deregistration. Canadian Tree Improvement Association Tree Seed Working Group News Bulletin 39: 9-11.

5.2.2 Western Redcedar Pollen Monitoring, Saanich Peninsula and Sechelt (SPU0212)

Joe Webber, ProSeed Consulting

Introduction

The Chief Forester's (CF) Standards for Seed Use in British Columbia have established the legal requirements for the use of tree seed on Crown land. For orchard seed lots, this includes effective population size (Ne) and genetic worth (GW). Calculations for estimating a seed lot's GW include female and male gamete contribution (including non-orchard sources of pollen and SMP).

Currently, only coastal Douglas-fir seed orchards must report contamination levels. However, other coastal orchard species (and interior lodgepole pine), including western redcedar, western hemlock, and Sitka spruce may also be at risk from contamination, but data to assess this risk is not available. We do not know the magnitude (pollen cloud density) and extent (duration of pollen flight) of any of these species. This report summarizes the first year results for monitoring western

redcedar pollen flight in two orchard locations: Saanich peninsula and Sechelt.

The Ministry of Forests' seven-day pollen monitors, mounted on rotating platforms at 3 m height, were used in each of two orchard locations; the Saanich peninsula and Sechelt. Two replicates were counted on each chart, and the total number of pollen grains for the two replicates were averaged for the two replicates and then divided by 44 mm^2 (the area of the chart counted) to represent the number of pollen grains (captured) per mm^2 for a 24-hour period.

Figures 20 and 21 show the mean western redcedar pollen catch for five stations on the Saanich peninsula and three stations at Sechelt, respectively.

In 2005, western redcedar pollen shed on the Saanich peninsula ranged from February 1 to February 28 with peak shedding occurring on February 19. It is likely that more pollen shed occurred later than February 28, but it was mixed with other species (predominately red alder pollen), which is the same the same size and shape. In terms of extent and magnitude, redcedar pollen shed on the Saanich peninsula is as great as or greater than Douglas-fir.

For the two western redcedar orchards on the Saanich peninsula (Western Forest Products, Saanichton and TimberWest, Mount Newton), the range of orchard receptivity was February 25 to March 9 and February 22 to March 11, respectively. This also corresponds to the maximum average daily pollen shed for contaminate pollen.

In 2005, western redcedar pollen shed at Sechelt ranged from February 10 to March 9 with peak shedding occurring on March 4. The magnitude and extent of other species (red alder) shed at Sechelt was less than on the Saanich peninsula. Douglas-fir pollen shed has not been monitored with the seven-day recorders at Sechelt, so we cannot make the same comparison as for the Saanich peninsula.

For the western redeedar orchards at Sechelt (Canfor), the range of orchard receptivity was February 18 to March 9. This also corresponds to the maximum average daily pollen shed for contaminate pollen.

Summary

At x100 magnification, it is possible to identify redcedar pollen, but it is very difficult to separate it from other species, in particular, red alder. Red alder is slightly smaller (20 to 25 microns in diameter), with pores. It looks very similar to redcedar at x100 magnification but with smaller orbicules. Positively distinguishing the two species at x400 is possible (glass slide and cover slip). However, distinguishing the two species at x400 on the Vaseline coated charts (no cover slip) is very difficult.

The pollen counts for western redeedar are as great as those observed for Douglas-fir on the Saanich peninsula, and the possibility of contamination exists. Whether this is a serious problem is not known. Any orchard clone sired with contaminated pollen could have a small reduction in selfing. However, the breeding value of contaminate pollen would at best be neutral, so any contamination will have a negative effect on higher-breeding-value clones.

This is the first attempt to monitor western redcedar pollen shed from indigenous stands surrounding the three orchard sites. The pollen monitoring technique used for monitoring Douglas-fir pollen in the orchards may not be applicable for western redcedar. Most orchards use dissecting microscopes with magnification ranges of x40 to x60. This is not suitable for monitoring western redcedar pollen. If orchard monitoring will be considered, then at least, a transmission microscope with a minimum of x100 magnification will be needed. Even then, distinguishing between redcedar pollen and other species, principally red alder, will be difficult, and the errors associated with the counts will be higher.



Figure 20. Mean daily western redeedar pollen catch (pollen grains/mm^2) for three stations at Sechelt. Black bars include other species of the same size and configuration (predominately red alder)



Figure 21. Mean daily western redcedar pollen catch (pollen grains/mm^2) for five stations on the Saanich peninsula. Black bars include other species of the same size and configuration (predominately red alder)

5.2.3 Damage Potential of Field Densities of *Leptoglossus occidentalis*

Ward Strong

The Western Conifer Seed Bug, *Leptoglossus occidentalis*, is the most serious pest in conifer seed orchards in the interior of BC Tests over the past six years have shown that it routinely reduces seedset by 10 to 50% and is capable of reducing seedset by up to 80%. *Leptoglossus* is one of the main factors contributing to the low-seedset problem in lodgepole pine orchards in the southern interior. It also feeds on white pine, Douglas-fir, spruce, hemlock, and larch, though the problem is less well characterized in these tree species.

Management of the seed bug depends on the use of broad-spectrum insecticides. Despite years of study, no other control methods, such as biological controls, trapping methods, or cultural techniques, have come to light. Appropriate use of insecticides requires two techniques: a method of estimating field densities of *Leptoglossus* (monitoring technique) and a method of relating these densities to damage potential (density-damage relationship). A visual-sample monitoring technique has been developed that consistently estimates seed bug densities. This project is designed to develop a density-damage relationship so that the damage potential of field densities of *Leptoglossus* can be estimated. 2004 was the final year of this three-year project

Methods

Leptoglossus densities were monitored weekly from early May through late July, 2004 at each pine orchard in the southern interior of BC Large pine orchards were divided in half, and each half was monitored separately. In total there were 15 orchards and half-orchards ("replicates"). Monitoring consisted of half-hour transects through the orchard on foot, visually examining cones and counting *Leptoglossus*. Monitoring in each orchard was also conducted in 2002 and 2003.

For each orchard replicate, accumulated *Leptoglossus*-days (LDs) were computed to estimate feeding pressure in that replicate. LDs were computed by taking the average of the current monitoring count and the previous monitoring count and multiplying it by the number of days between counts. For example, if the count on June 10 was 12 *Leptoglossus*, and the count on June 17 was six *Leptoglossus*, LDs accumulated in that period was $(12 + 6)/2 \times (17 - 10) = 63$ LDs.

In each orchard replicate, feeding damage was assessed with insect exclusion bags. On April 28, 2004, exclusion bags were placed over second-year cones to protect them from *Leptoglossus* feeding. On July 30, bagged cones and nearby unbagged cones (exposed to *Leptoglossus* feeding) were collected. Seeds were extracted by the boil, bake, and shake method, and seedset (filled seeds per cone) was computed for each sample. Seedset loss (the difference in seedset between exposed and bagged cones) was a measure of feeding damage. Seedset loss was also calculated for 2002 and 2003 data.

Feeding damage was then graphed against LDs to indicate the relationship between monitored densities of *Leptoglossus* and feeding damage. A regression line was generated to describe the relationship mathematically. A close relationship is indicated by a high r^2 value of the regression. The closer the relationship between LDs and damage, the more reliably we can estimate damage from monitored densities.

Results

The bagged and unbagged treatments have been harvested from each orchard replicate, and the seeds extracted and X-rayed. The filled and empty seeds were counted. Seed loss due to *Leptoglossus* for each orchard replicate was then calculated. For the rest of this report, results from all years of the study will be discussed together (2002, 2003, and 2004 combined).

Scatterplots of feeding damage as a function of *Leptoglossus*-days were generated. Figure 22 shows the reduction in seedset (filled seeds per cone) as a function of LDs. Lepto days were calculated in three ways:

A. Total LDs: all *Leptoglossus* were counted equally (females, males, and nymphs)

B. Female LDs: Only the females were included. This is because the females feed much more than males or nymphs and cause the greatest amount of damage

C. Low-male LDs: Males and nymphs feed about one-third as much as females, so LDs were calculated by counting each female as one lepto and each male or nymph, as 0.3 lepto.

On the scatterplots, each point represents one orchard replicate in one year. Female LDs resulted in the poorest relationship, and low-male LDs resulted in the

best relationship, as identified by the higher r^2 value for the low-male plot.

The r^2 value for low-male LDs is still low, however: $r^2 = 0.1134$ (Figure 22C), which is a relatively poor relationship. Several other factors contribute to seedset reduction, such as the crop size. A certain number of *Leptoglossus* will reduce the FSPC in a small crop more than in a large crop. Crop size estimates (by the orchardists) were used to generate an estimated number of seeds lost for each orchard replicate in each year. Seeds lost were then plotted against LDs; these are shown in Figure 23.

The relationship is much tighter than with the FSPC reduction data. This means predictions of damage can be made with more accuracy. Total LDs and low-male LDs had about the same r^2 values, at about 0.18. These figures can be used to estimate seed loss due to varying densities of *Leptoglossus*. For example, using the total LD graph (Figure 23A), we can estimate that if we have 600 LDs of feeding pressure, we will lose approximately 1.25 million seeds from that orchard.

Because of the poor relationship (low r^2 value), this estimate is not very accurate though. Other factors that contribute to variability in the density-damage relationship fall into three categories:

A. Orchard influences – e.g., crop size, orchard size, orchard age, tree size, spacing

B. Monitoring influences – factors that contribute to monitoring results: e.g., temperature, cloud cover, wind, time of day

C. Seasonality – In a related experiment in 2004, I found that feeding in spring caused far more damage than feeding in summer.

These factors will be incorporated into the density-damage relationship as data analysis proceeds. The orchard influences are known and easily incorporated. The monitoring influences and seasonality have now been characterized and can also be included.

Once an accurate density-damage relationship has been described, we will be able to make more accurate predictions about seed loss due to monitored numbers of *Leptoglossus*. This will help us make well-informed decisions about managing *Leptoglossus*. Pesticide spray thresholds will be the main decision-making tool. We might be able to say, for example, that "x number of *Leptoglossus* found in our half-hour search indicates a potential seed loss of y%, and therefore a spray is not warranted." In this way, pesticide applications will be minimized and will be used only when and where necessary.


A. Total Lepto Days vs Seedset Reduction
B. Female Lepto Days vs Seedset Reduction
C. Low Male Lepto Days vs Seedset Reduction

Figure 22.



Figure 23.

5.2.4 Douglas-fir and Western Redcedar Cone and Seed Insect Pest Monitoring Services to Coastal Conifer Seed Orchard Managers.

Robb Bennett

Douglas-fir cone gall midge and western redcedar cone midge are the two most important insect pests in coastal British Columbia conifer seed orchards.

Population levels of both these insects are routinely monitored each spring to determine damage potential and whether or not control options need to be used. Orchard managers are now responsible for monitoring these insects on their sites. Formerly, Ministry of Forests Seed Pest Management staff provided this service. This project was initiated to ease the transition process for coastal seed orchard managers.

In spring 2004, conelet samples from eight of eleven coastal Douglas-fir seed orchards at four sites were assessed by Applied Forest Science Ltd. for levels of Douglas-fir cone gall midge. Reports of results (Figure 24), including pest management recommendations, were sent electronically to respective orchard managers within 24 hours of receipt of samples. The remaining orchards were either unproductive in 2004 or were being monitored by on-site staff.

Redcedar conelet assessments were done in-house in March 2005 by orchard managers, following successful training in monitoring techniques provided by Seed Pest Management staff.

Orchard	ISCC	Orchard	ISSC
116	11.2	166	4.6
120	9.3	169	3.2
149	0.9	177	14.4
162	0.4	181	6.5

Figure 24. Average infested scales per conelet counts (ISCC) for eight of eleven coastal BC Douglas-fir seed orchards. Default "treatment threshold" (at or above which control measures are usually warranted): ISSC = 5.0 6

5.2.5 Estimation of Deleterious Effects of Different Levels of Inbreeding on Western Hemlock Seed Orchard Production

District Hub

In forward-selected advanced-generation seed orchards, and even in first-generation orchards where SMP is used, there is likely to be some reduced seed set due to inbreeding. The size of these effects for western hemlock is unknown, except for selfing. Previous studies have shown that the propensity of western hemlock orchard ramets to self-pollinate shows a strong family effect. This factor is particularly important in clonal row orchards. Once problem clones are identified, managers can avoid placing ramets of these clones adjacent to each other, or they can employ some form of pollen management: e.g., supplemental mass pollination.

Even low-level inbreeding has been shown to reduce seed production. It is logical to assume that there will be problems, but the size of these effects can vary considerably even within a botanical family such as the *Pinaceae*. It is therefore advisable to investigate a low-level selfing for a species before designing advanced-generation orchards since impacts on production should be taken into account. Minimizing selfing in orchard seed will ensure that inbreeding does not compromise the growth potential of high-gain lots and that there is minimal reduction in seed set.

In this project, 48 high-breeding-value orchard clones were tested. The clones were then ranked from the lowest to the highest level of selfing, based on their propensity to self. The final results are presented in Figure 25.

	Clone	% Full Seed Set[a]	% Full Seed Filled[a]	Level of Selfing
1	356	1	45	very low
2	1084	2	36	very low
3	376	2	39	very low
4	748	2	39	very low
5	2049	3	71	very low
6	122	3	68	very low
7	2166	3	53	very low
8	489	3	57	very low
9	756	4	71	very low
10	61	4	55	very low
11	497	4	62	very low
12	4492	5	58	very low
13	4230	5	65	very low
14	2238	6	55	very low
15	1002	7	83	very low
16	2575	8	76	very low
17	810	8	67	very low
18	440	8	56	very low
19	196	7	98	very low
20	31	6	48	very low
21	2252	8	60	very low
22	248	9	57	low
23	445	9	55	low
24	2520	13	61	low
25	462	13	53	low
26	230	12	60	low
27	569	15	75	low
28	48	15	69	low
29	491	15	69	low
30	120	13	51	low
31	2007	17	89	moderate
32	268	17	75	moderate
33	2460	18	77	moderate
34	60	16	96	moderate
35	459	21	77	moderate
36	140	19	91	moderate
37	1014	21	68	moderate
38	457	27	71	high
39	49	25	56	high
40	1203	32	68	high
41	2465	30	76	high
42	169	33	73	high
43	1203	45	68	high
44	74	48	85	high
45	4091	53	79	high

[a] Full empty seed (FES)

Figure 25. Propensity to self. Ranking of the clones from the lowest to the highest level of selfing.

1.2.6 Determination of Selfing Rates in a Top-pruned Interior Spruce Seed Orchard (SPU #30)

Michael Stoehr and Hege Kiehl

Background

Interior spruce orchards are routinely top pruned for more efficient orchard management and increased safety during cone harvest. However, it is prudent to determine if this management practice affects seed yield and seed quality. Effectively, top pruning brings the male and female reproductive buds closer together, creating a wider zone of overlap which could potentially result in higher selfing rates. To assess actual selfing, this orchard was genotyped using chloroplast SSR markers (last year's project) and cones from two candidate clones were collected from the lower, the middle, and upper sections of the crown.

Activity

Cones from two ramets each of ten clones that were identified as unique, based on our SSR analysis of last year, were collected in July 2004 from the lower, the middle, and the upper part of the crown. Cones were after-ripened in a growth chamber prior to seed extraction. 80 seeds were extracted from each cone and x-rayed to determine the proportion of filled seed.

Results

The molecular SSR analysis was not finished, and only the seed yield data are reported. Averaged over all tested clones (two ramets each), the number of seed per cone was lowest in the lower part of the crown, intermediate in the mid-crown, and highest in the upper crowns. The same trend held for the number of filled seed. The proportion of filled seed per cone ranged from 34% to 44% in cones growing in the lower and upper crown, respectively. In the absence of SSR data, it is difficult to interpret the results because the lower number of filled seed in cones growing in the lower part of the crown may be due to the lower vigour of lower branches and/or selfing.



	Lower	Mid	Upper
Total Seed	114	137	163
Filled Seed/%	39/ 34	55/ 41	69/ 44

Figure 16. Effects of cone position on seed yield per cone in a top-pruned interior spruce seed orchard

1.2.7 Development of Pollen Management Guidelines for Yellow-cedar

Dave Kolotelo

Poor pollen quality at pollination time may be one of the principal factors responsible for the failure of low-elevation orchards to produce sufficient quantities of viable seed. In a pollen quality study, significant differences in pollen viability (pollen germination) were observed between populations characterized by distinct environmental differences. There was a significant reduction in pollen development and in corresponding pollen quality as the elevation of the rearing sites decreased (Figure 17).

All previous assessments of pollen quality were based entirely on in vitro testing, using pollen germination. These assessments should be interpreted only as an indication of pollen fertility. In the continuation of this project, the actual fertility of low-elevation males is being confirmed through control pollinations of females in natural stands. The level of fertility of low-elevation females is also being examined through control pollinations using high-quality pollen collected from natural stands.

Results to date indicate that low-elevation yellow-cedar seed orchards are capable of producing viable seed. The study further shows that both low-elevation males and low-elevation females can produce viable seed

(Figure 4B). For example, in a high-elevation stand, one observation made produced a similar percentage of filled seed and a similar number of seeds per cone as high-elevation trees. Comparable results were observed at low elevations where there was no insignificant difference in the percentage of filled seed or the number of filled seeds per cone between matings with high-elevation and low-elevation pollen.

Even though females were able to produce filled seeds at low elevations, the percentage of filled seed and the number of filled seeds per cone was significantly lower than at high elevations for the same mating types. However, when open-pollinated cones were collected at low elevations from the same trees used for control pollination, they all produced empty seed.

These data do not contrast to earlier reports that low-elevation seed orchards on Vancouver Island are not capable of producing viable seed. One possible answer to this discrepancy lies in the pollination method employed. Open pollination was in effect in the early seed orchards at low elevations. Similarly, when open-pollinated cones were collected in the current study at low elevations, they also produced empty seed. However, when control pollination was used in the current study, viable seed was produced. There was an abundance of pollen on the mated trees and on surrounding trees during open pollinations. This suggests that there may be several unknown and untested factors responsible for this failure.

Although there is no significant difference in filled seed between high x high and high x low mating types at high elevations, and between low x high and low x low at low elevation, it is almost significant and probably biologically significant. There was a lot of trees by trees delay variation, and the number of trees at each location was reduced from ten to six trees at high elevations and from ten to eight trees at low elevations. Difficulties with environmental conditions encountered, especially at high elevations, reduced the number of mated trees and also introduced several other factors that may have influenced, to a certain degree, the results of the trial (see "Results" above). These problems are being dealt with in sub-project 3.



Figure 4. Relationship between yellow cone pollen ability to pollinate and susceptibility to cone insects and cone crop losses.

Figure 4B. Yellow cone pollination study. Filled seed for two mating types.

[illegible]

[illegible]

Female reproductive [illegible] has adapted to cold climates and short growing seasons may be explaining different seed growth variations and [illegible] range growing seasons. The highest [illegible] yellow pollen in cool growing seasons is cold climates. For example, [illegible] in temperate conditions, can find considerable [illegible] range in [illegible] [illegible] [illegible]

[redacted]

[redacted]



[redacted]

[redacted]

[redacted]

[redacted]

[redacted]

[redacted]

methods. Leaving the trees unmanaged is definitely not an option in a micro orchard setting. Because space is limited, it is essential to manage the growth of each ramet so that it does not expand out of its allotted area.

The three crown management methods deliver reasonably close results. Operational-style pruning seems to yield the highest number of cones. This treatment is also the most efficient to implement because it does not require any labour intensive branch training.

Conclusion

We have been testing our current crown management techniques in orchard #505 for four field seasons. Because our survey results are based on three years' worth of inconsistent flower crop data, we are reluctant to make any firm recommendations.

At this point it appears that operational-style crown pruning may be the most effective method to apply to the interior spruce micro orchard.

In 2005 we will continue to refine our techniques in both orchards and collect additional crop data.



Figure 37. No crown management effect on crop

5.2.10 Testing the Effects of Upgraded Irrigation Delivery Systems in NE Mid-interior Spruce Seed Orchard #305 and Interior Lodgepole Pine Seed Orchard #307

Gary Comps, Kalamalka Seed Orchards

Introduction

In 2003 OTIP funded revisions to the irrigation delivery systems in portions of interior spruce orchard #305 (SPU6x12) and interior lodgepole pine orchard #307 (SPU6718). Our 2005 study is a logical continuation of these projects. We are interested in learning how the upgraded irrigation system impacts flower production and ramet health. This is intended to be an ongoing study that tracks and compares ramet performance throughout several growing seasons.

Background

Orchard #305 was established at Kalamalka in 1981. Orchard #307 was planted in 1986. Since their inception these orchards have been watered with a drip-style irrigation system. The trickle system seemed to work well enough when the orchards were young, but as the trees matured we began to question the efficiency of this type of irrigation delivery. Delivering water to the base of each tree limits root growth to the zone wetted by the emitter. As the tree grows larger this limited zone may constrict root volume expansion to the point where the roots cannot service the developing crown. This imbalance could have a negative effect on ramet health and might ultimately lead to decreased seed production.

A portion of each orchard has had its irrigation system converted to broadcast delivery as planned. We began watering operationally with the converted system in 2004.

The portion of the orchards serviced by the broadcast delivery system will be compared to the portion of the orchards irrigated with the drip system. These observations will allow us to determine how the different irrigation delivery systems affect ramet performance.

Activities

In the summer of 2004 permanent sample trees were chosen in each orchard (96 trees in orchard #305 and 84 trees in orchard #307). The sample trees were chosen

in pairs based on establishment date: one tree is in the area irrigated with the micro sprinklers and the other is in the drip-irrigated area.

Three major branches on each sample tree were chosen at random and marked with metal tags. Shoot elongation on each branch was measured during 2004. We also measured stem diameter below all the branches on each sample tree. A survey was conducted to observe flower production. We will revisit these trees on an annual basis to repeat these observations.

Cone samples representing the two different irrigation regimes were harvested from each orchard. Seed has been extracted from these cones to compare seed production. Foliar samples were also collected to analyse nutrients.

Results

The new irrigation regime has only been in effect for one growing season and may not yet have made a large impact on ramet performance. None of the shoot or diameter measurements indicated any differences between the two irrigation treatments at this point. The data collected in 2004 will serve as a baseline for future observations.

Seed was extracted from the cone samples and X-rayed. At this point there is no significant difference in filled seed per cone or total seed per cone between the two treatments.

Nutrient uptake was one area where broadcast irrigation seemed to make a difference. When the foliar samples were processed, higher levels of aluminium, boron, calcium, manganese, nitrogen, and phosphorus were observed in the samples collected from the portions of the orchards watered with the broadcast delivery system.

Conclusion

We will continue to monitor the permanent sample trees on an annual basis. A set of comprehensive baseline observations will be collected over the course of several growing seasons. Comparing these results will allow us to determine the effectiveness of broadcast delivery and make recommendations to other orchard managers considering irrigation system upgrades.

5.2.11 Clonal Verification of a Spruce Weevil Orchard (#172) Using DNA Markers (SPU 0611)

Craig Newton and Tam Vo

Clonal propagation (CP), either by rooted cuttings, grafting, or more recently through somatic embryogenesis, is an essential component of some breeding and tree improvement programs. Through CP, individual trees with superior traits (genotypes) can be multiplied to create sufficient numbers of propagules for field trials, seed orchard establishment, and any other application for which identical genetic backgrounds are needed. All these applications are dependent on the fidelity of clonal propagation techniques and the labelling systems in place to track them through different seed orchards and test plots.

To test the fidelity of clonal propagation and identification management, a high-gain Sitka spruce weevil resistant seed orchard (#172, Western Forest Products) was tested using nuclear micro-satellite markers. This class of DNA marker is equivalent to those used in human forensics cases and can produce multi-locus genotypes that are unique to every individual tree. Therefore, for each orchard parental clone, the original ortet and all ramets produced from that ortet should share identical genotypes if clonal propagation and labelling systems are working as predicted.

In this study, 144 ramets representing 15 different clones (3-15 ramets per clone) were analysed with four nuclear micro-satellite loci. A total of 6/144 ramets (4%) were different at all four loci and therefore represent cases of non-clonal ramets. Figure 38 shows examples of non-clonal ramets identified with one of the four micro-satellite markers (SSR44). The origin of some of the verified non-clonal ramets was also determined – two of the 887 ramets were identical to clone 945, and two non-clonal 1056 ramets are identical to clone 1253. Ortet material from clones 887, 1056, and 1253 was also tested and found to be identical to the majority ramet genotype.

In conclusion, clonal fidelity in orchard #172 was 96%, and most (4/6) of the non-clonal ramets appear to be due to mislabelling rather than biological events such as rootstock over growth. Ortet material from three of the 15 families tested was also identical and indicate overall that clonal fidelity in orchard establishment is high.



Figure 38. Clonal verification in Sitka spruce using micro-satellite markers

5.2.12 Enhancing Seed Production in North Okanagan Lodgepole Pine Seed Orchards (SPU 0719)

Chris Walsh and Joe Webber

This report summarizes the fifth year of the results from studying the effect of culture (irrigation and crown cooling) on seed set in the Kalamalka lodgepole pine seed orchard #307. After five years, results clearly show that operational quantities of seed (15 filled seed per cone) can be obtained if aggressive insect protection measures are taken.

Results from years 2000-2004 show no significant effect of enhanced culture on seed yields, although cone mass and seed weights were improved slightly. There still remains concern that hot, dry weather during the pollination period could result in poor pollen uptake and subsequently lower seed yields. Accordingly, objectives for 2004/05 focused only on treatment (irrigation and crown cooling) effects applied during the pre-pollination (mid-April to mid-May) and pollination (mid- to late-May) periods. Both irrigation and mist systems were activated in early April. Mist systems were deactivated after pollination (late May) and irrigation systems were deactivated at cone collection (early August).

Treatment effects on seed yields were measured (total seed per cone, filled seed per cone, and seed weight), cone mass (first-year conelets and second-year cone dry weights), and cone numbers. Meteorological data (temperature and relative humidity) were also recorded and expressed as heat sums and vapour pressure deficit.

Figures 39 and 40 show the meteorological data for the two test periods, pre-pollination and pollination, respectively. Included are sums of mean hourly temperature and vapour pressure deficit, as well as sums where the mean hourly temperatures exceeded 24°C (and the number of hours) or where vapour pressure deficit exceeded 2 (and the number of hours). Vapour pressure deficits exceeding 2 are associated with drought conditions. Precipitation is also shown.

The weather during the pre-pollination period was warmer than usual although not drier than previous years. However, during the pollination period, temperatures remained relatively warm while precipitation was about three times normal. After the pollination period, hot dry weather resumed.

Figure 41 shows the mean total seed per cone (TSPC), filled seed per cone (FSPC), and seed weight for cones from the control block (no treatments) with and without insect protection for the last five years.

Figure 42 shows the mean cone number per major whorl branch (pollen bud clusters, first-year conelets, and second-year cones), and cone weights (first-year conelets and second-year cones) for cones from the control block (no treatment) for the last five years.

Meteorological data in Figure 40 show that the pre-pollination period was warmer than usual but also considerably wetter. We were not able to test adequately the study objectives of hot dry weather during pollination on pollen uptake.

However, the data for seed yields (Figure 41) and cone numbers and weights (Figure 42) show some interesting trends. These data are for the control block only. This block was untreated (no irrigation or crown cooling). Although data for the treatment blocks do not substantially differ from those for the control block, the data for the crown cooling and irrigation blocks are available in the final report for this project.

The most significant loss of seed yields continues to be attributed to insect (*Leptoglossus*) damage. Over the five-year period, mean filled seed per cone for protected and unprotected cones was 22.8 and 15.8, respectively. Similar losses in total seed per cone were also observed (32.9 versus 26.9, respectively). Seed weight was not affected by insect feeding.

There is concern that seed cone numbers have steadily decreased over the study period. Pollen cone clusters, however, have increased. This trend is likely the result of two important factors. First, the same major whorl branches on each of the sampling trees were used for the five-year period. Cone numbers and seed yields are significantly greater in the upper crown. As the trees grew larger, the sampling branches became lower in the crown. Branch vigour became lower and more suited to pollen production. Second, weather conditions during the cone differentiation period for seed cones (September) has been cool and moist in the last three years. However, first-year conelet and second-year cone weights have remained reasonably consistent over the five-year period.

Our conclusions for last five years remain the same. Controlling *Leptoglossus* is the single most important management activity for lodgepole pine seed orchards in the north Okanagan. Cultural treatments (irrigation) do provide a modest increase to seed yields and cone size, but the beneficial effect of misting during the pollination period has yet to be determined. While the increased irrigation capacity provided only a slight improvement to seed yields and cone size (relative to the previous drip system), it seems prudent to maintain these systems knowing that the root volume for the matures trees is larger and the trees have a better chance of surviving extreme temperatures and drought when they do occur.

	2000	2001	2002	2003	2004
Tsum	na	4740	3673	4400	5365
Tsum>24	na	299	74	48	456
T>24 (hrs)	na	12	3	2	18
VPDsum	na	518	503	414	547
VPDsum>2	na	39	45	28	67
VPD>2 (hrs)	na	17	20	13	29
precipitation (mm)	na	16.9	13.9	73.4	60.4

Tsum=sum mean hourly T-5°C
Tsum24=sum mean hourly T>24°C
VPDsum=sum mean hourly VPD
VPDsum2=sum mean hourly VPD>2

Figure 39. Temperature (T) and vapour pressure deficits (VPD) sums and precipitation (mm) for the period of April 15 to May 15 for the study period of 2000 to 2004

	2000	2001	2002	2003	2004
Tsum	3466	4100	3603	3771	3631
Tsum>24	104	1515	304	48	578
T>24 (hrs)	4	54	12	2	18
VPDsum	313	424	283	414	283
VPDsum>2	23	163	50	54	67
VPD>2 (hrs)	10	56	22	22	29
precipitation (mm)	na	20.1	38.5	38.6	143.8

Tsum=sum mean hourly T-5°C
Tsum24=sum mean hourly T>24°C
VPDsum=sum mean hourly VPD
VPDsum2=sum mean hourly VPD>2

Figure 40. Temperature (T) and vapour pressure deficit (VPD) sums and precipitation (mm) for the period of May 16 to May 31 for the study period of 2000 to 2004

	2000		2001		2002		2003		2004	
	Bag	NoBag	Bag	NoBag	Bag	NoBag	Bag	NoBag	Bag	NoBag
TSPC	na	22.4	36.9	30.6	33.3	27.5	31.0	31.1	30.4	22.9
FSPC	na	9.7	25.5	18.8	24.3	16.7	21.4	19.8	20.1	13.9
Seed Weight (mg)	na	4.1	3.2	3.0	4.2	4.1	4.3	4.2	4.4	4.4

Figure 41. Mean total seed per cone (TSPC), filled seed per cone (FSPC) and seed weight for cones with (Bag) and without (NoBag) insect protection for the study period of 2000 to 2004.

	2000	2001	2002	2003	2004
Cone Numbers					
Pollen Clusters	37.6	46.7	61.1	70.6	74.6
Conelets	16.8	16.7	12.7	8.8	4.9
Cones	7.9	15.4	13.4	10.5	6.9
Cone Weights (g)					
Conelets	0.229	0.195	0.182	0.188	0.209
Cones	6.54	6.86	6.19	6.93	6.10

Figure 42. Mean cone number per major whorl branch (pollen cone clusters, first-year conelets, and second-year cones), and cone weights (first-year conelets and second-year cones) for the study period of 2000 to 2004

5.2.13 Chemical Control Techniques Against *Dioryctria abietivorella* in Interior Fd and Pw Orchards.

Ward Strong

Douglas-fir, spruce, and white pine orchards in the interior are now producing harvestable crops of cones. It has become apparent that *Dioryctria abietivorella* is a major pest of these crops. *Dioryctria* is a moth whose larvae bore inside cones, feeding on cone tissue and seeds. Losses in 2003 exceeded 40% in some crops.

This OTIP project was designed to meet the immediate need for control measures to protect crops until the *Dioryctria* are better understood. The project was to consist of two components: bagging studies and pesticide trials. Due to factors beyond my control, I was able to conduct only part of the planned pesticide trials. These are reported here.

Dimethoate is a systemic organophosphate registered against cone and seed pests in Douglas-fir. When sprayed onto a tree, the pesticide is translocated inside the cones, where it controls pests that feed on cone and seed tissue. Two experiments assessed its efficacy against *Dioryctria*: one with a backpack sprayer on individual trees and one with an orchard air-blast sprayer on large blocks of orchards. In each, dimethoate was applied once in late April or twice in late April and mid-June. The late April applications were at high and low rates, and the later application was at the low rate only. Cone damage was checked in mid-June and late July; damage on treated trees was compared with untreated checks.

Results are shown in the figures below. In both trials, the early spray reduced damage significantly on the mid-June check, but on the trees with early sprays only, damage was showing up by mid-July. Trees sprayed twice still had very low damage by 14 July.

It appears that *Dioryctria* moths continued to lay eggs throughout the experimental period, requiring active insecticide within the cones to kill hatching larvae. The early application provided control for perhaps six weeks, after which control declined, resulting in a higher percentage of infested cones by mid-July. When two applications were made, control continued through mid-July. Although no counts were made after this date, it was apparent at harvest that damage had started to increase in the trees with two applications, suggesting about six weeks of control from the second application as well.

Perhaps the best use of this chemical would be to wait until damage is first noticed, sometime in late May, before making the first application, then make a second application in six weeks, which should carry protection through the harvest period. However, the pesticide label allows an application only immediately following cone receptivity, thus the label must be amended before the desired use is possible. In 2005 we will be investigating an URMULE minor-use label expansion for this purpose.

Dimethoate	22 June	14 July
5% April 19	3%	39%
2% April 19	9%	36%
5% + 2% Jun 20	6%	4%
No spray	31%	60%

Figure 43. Orchard block trials. Percentage of cones infested with *Dioryctria* in blocks sprayed with different concentrations of Dimethoate 480E

Dimethoate	15 June	21 July
2.5% April 22	0%	38%
1% April 22	1%	57%
2.5% + 1% Jun 15	1%	7%
No spray	44%	90%

Figure 44. Individual tree trials. Percentage of cones infested with *Dioryctria* on individual trees sprayed with different concentrations of Dimethoate 480E

Plate 21. *Dioryctria abietivorella* adult

Plate 22. *Dioryctria* damage on Douglas-fir cones

Where We Work

Tree Improvement Program 2004/05





6.0 Extension and Communications

6.1 Extension Technical Advisory Committee (ETAC Subprogram)

Dave Douglas

Annual ETAC meeting

The annual ETAC business meeting was held in Kamloops on November 4, 2004. Kathie Swift of FORREX gave a presentation on Extension Planning Workshops.

Extension Notes

Extension notes initiated over the past few years are now completed and available on the FGC website at http://www.fgcouncil.bc.ca/index.html:

Extension Note 3, Benefits of Using Selected Reforestation Materials, Carlos and Jack Woods, January 2005

Extension Note 4, The Reproductive Biology of Western White Pine, April 2004 (100 copies of Extension Note 4 were printed.)

Extension Note 5, Environmental Effects on Yellow-cedar Pollen Quality, Dietrich Hak, April 2004

Extension Note 6, Applications of DNA Markers in BC Tree Improvement Programs, Craig Newton, April 2004

Extension notes in production this year are:

The Reproductive Biology of Lodgepole Pine

Use of DNA Markers to Evaluate SNP Efficacy in Lodgepole Pine

Cone Induction in Western Red-cedar Seed Orchards

Somatic Embryogenesis Project Report

ETAC Client Survey

The Forest Genetics Council Tree Improvement and Genetics information needs survey was developed by FORREX and delivered in January 2005 through a web-based program called zoomerang. SPAR, BCTS, woodlot owners, BCMoF, consultants, and other industry contacts comprised a client contact list of 316 persons. Results of the survey will be presented to the Forest Genetics Council and published. An Extension Planning Workshop will be the next step in developing an Extension Plan for FGC through ETAC.

Tree Improvement in BC brochure

This brochure is currently being revised and updated.

Northwest Forest Genetics Tour

A forest genetics tour was held October 6, 2004 to view some of the provenance/progeny trials in the Kitwanga/Kispiox area of BC. Issues that were discussed ranged from seed "transfer" guidelines, diseases of Lodgepole pine, Chief Forester's Standards for Seed Use, genetic gains, and genetics relative to site productivity and global warming.

Stops were at Will Cook (hybrid spruce trials and results); Kitwancool Lake (western hemlock, yellow-cedar, western redcedar testing, and *Dothistroma* damage in Lodgepole pine); Kitwanga (species trials); and Helen Lake (hybrid spruce and western hemlock trials). Alvin Yanchuk, John King, Greg O'Neill, and Charlie Cartwright, Research Branch; Alex Woods, regional pathologist at Smithers, and Brian Barber, Tree Improvement Branch, made presentations. Representatives from BCTS, industry, BCMoF, and forestry consultants participated.



Figure 41. DF 702.04 and 1972 trial locations



Plates 15 and 16. Photos from the forest gazebo near Kabili-Kindoki in June 2004 in close vicinity of the primatological programs made in the Kisangani/Yoko area of DR.

2.0 Seed Pest Management

2.1 Operational Assay Program for Fungi Attacking Conifer Seed in BC

[illegible]

The BC Ministry of Forests Tree Seed Centre leads this project, with collaboration from Applied Forest Science Ltd. Now in its second year, the project's objectives are to:

- continue to perform fungal assays on seed lots from conifer species known to be at high risk to infection by a particular pathogen and
- make this information available to seed users through the Seed Planning and Registry System (SPAR).

The following fungal assays were performed in 2006/07:

- Cylindrocarpon (trigger) – 48 seed lots
- Fusarium spp. – 99 seed lots
- Rhizosphaera spp. – 48 seed lots

The results of these assays were posted on SPAR in preparation for the 2007 seed lot ordering deposits.

2.2 Identification of an Effective Sex Pheromone Lure for the Fir Coneworm, *Dioryctria abietivorella*, and Demonstration of its Efficacy in Seed Orchards (SPH 06-02)

[illegible]

Canadian Forest Service and University of California (Riverside) researchers led the second year of this project with collaboration from the BC Ministry of Forests. Primary objectives of this work are to:

- identify the sex pheromone of the fir coneworm and
- develop this pheromone as a tool for sampling fir coneworm populations and monitoring coneworm phenology in seed orchards.

Key significant progress was made in 2006/07. Problems identified in 2005/06 field and lab work with pheromone components were resolved. Interactions (at dosages showed that the standard commercial preparation of one of the pheromone components was contaminated. Lab bioassays and field trials allowed identification of a previously unknown component of the fir coneworm sex pheromone. The new component was successfully synthesized and subsequently field tested in conjunction with the two previously identified and now determined to be only components of the sex pheromone. For the first time, large numbers of male coneworm moths were captured. Results indicated that one of the previously identified components is actually inhibiting and that a blended ratio of the remaining two components is critical. In 2006/07, the final year of this project, simple synthesis techniques for the pheromone components will be developed, and field trials with blending the pheromone blend for operational field use.

2.3 Identification of Seed Chalcids Infesting Seeds of BC Conifers (SPH 06-03)

[illegible]

2006/07 marked the second year of forest operations' support for this long-standing international collaborative project led by the Canadian Forest Service (CFS) and the National Museum of Resources Agroecologique (INRA), France. The BC Ministry of Forests (BCFP), the Tree Seed Centre and the Seed Pest Management groups and Nature Trust Propagation Unit are major collaborators. The long-term objectives are to:

- clarify the identity and host range of the North American species of seed insects in genera, and of those exploiting conifer seed in BC in particular, and
- determine the identity of cues (host) forming kits for parasitoid insects attacking seed wasps.

The 2006 growing season was poor, with crops in most natural stands across the province. Addressing the

[redacted]

[redacted]

[redacted]

Appendix 1 FY21 Seed Planning Limit

No.	Seed Planning Unit (SPU)	Name	Unit (Kilogram (kg) or Litre (l) or Number (no.))	Duration (start date - end date)		Lead time (days)	
				Min	Max	Min	Max



Appendix C: Tree Species

[redacted]	[redacted]	[redacted]
[redacted]	[redacted]	[redacted]
[redacted]	[redacted]	[redacted]
[redacted]	[redacted]	[redacted]

Appendix 3 Discussions by Tree Species

CONIFERS	ABBREVIATION	PAGE LOCATIONS
Western redcedar	Cw	7, 18, 21, 22, 32, 37
Yellow-cedar	Yc	9, 19, 21, 22, 39-42
Douglas-fir	Fdc	5, 18, 21, 22, 23, 31, 37, 48
Interior Douglas-fir	Fdi	10, 24, 25, 26, 43
Amabilis fir	Ba	22
Western hemlock	Hw	6, 18, 22, 31, 38
Western larch	Lw	36, 24
Lodgepole pine	Pli	12-13, 24, 25, 26, 27-29, 35-37, 44-45, 46-47
Western white pine	Pw	8, 14, 21, 22, 24, 29, 31, 48
Sitka spruce	Ss	5, 18, 45-46
Spruce hybrid (interior spruce/Ss)	Se	11-12, 24, 25, 27, 29, 38, 43
Ponderosa pine	Py	14-15
HARDWOODS		
Hybrid poplars	Ac	
Paper birch	Ep	
PEST		
Caloscypha fulgens		54
Fusarium		54
Sirococcus		54
Diorygtria abietivorella		54, 48-49
Seed chalcids		54-55
Leptoglossus occidentalis		35-37, 55

Appendix 4 Authors and Species Topic

AUTHOR	SPECIES/DISCUSSION	PAGE
Aitken, Sally	Gene Conservation	3-4
Ashley, Valerie	Fdi, Lw, Se	11-12, 16
Bennett, Robb	Cone and Seed Pests	32, 37, 54-55
Berger, Vicky	Pli	12-13, 14-15
Bird, Keith	Fdc	5
Brown, Patti	Fdc, Hw, Pw, Yc, Cw	21
Carlson, Mike	Pli	12-13, 14-15
Cartwright, Charlie	Hw, Ss	6-7
Cox, Keith	Pw, Se	14, 15, 29-30
Crowder, Tim	Fdc, Pw, Ba, Hw	21-22
Douglas, Diane	Extension	52-53
Ferguson, Craig	Cw, Yc	7-9
Glampa, Gary	Se, Lw, Pli	43-45
Graham, Hilary	Pli, Fdi	26-27
Hak, Oldrich	Hw, Yc	38, 39-42
Hoege, Bonnie	Fdi, Se, Lw	11-12, 16
Hollefreund, Clint	Fdc	5
Hunt, Rich	Pw	9-10
Jaquish, B.	Fdi, Se, Lw	11-12, 16
King, John	Ss, Se, Pw	5-6, 9-10
Kolotelo, David	Cones and Pests	54
Lee, Tim	Se, Pli, Fdi	25
Lohr, Carolyn	Fdc, Hw, Pw	31
McAuley, Leslie	SPAR	2
Mehl, Helga	Se	39
Murphy, John	Pli	12-13, 14-15
Nicholson, George	Pli, Se	27
Newton, Craig	Se, Lw	45-46
Painter, Roger	Program Management/Extension	17-18
Phillips, Giselle	Fdi, Se, Lw	11-12, 16
Piepet, Greg	Pli, Se	27
Ponsford, Dave	Ss, Pw	5-6, 9-10
Rudolph, Dan	Pw, Fdc	22-23
Russell, John	Cw, Yc	7-9
Stoehr, Michael	Fdc, Pli, Se	5
Strong, Ward	Leptoglossus	35-37, 48-49
Van Niejenhuis, Annette	Cw, Fdc, Hw, Ss	18-21
Wagner, Rita	Pli, Se	27-29
Walsh, Chris	Pli, Se, Lw, Pw	14-15, 23-24, 46-47
Webber, Joe	Pli	32-34, 46-47
Wigmore, Bevin	Cw	21-22
Woods, Jack	Program Management/SelectSeed	1

Appendix 5 Author Contact List

CONTRIBUTOR	AFFILIATION	TELEPHONE NUMBER
Aitken, Sally	UBC	604-822-6020
Ashley, Valerie	MoF	250-260-4753
Berger, Vicky	MoF	250-260-4767
Bennett, Robb	MoF	250-652-6593
Bird, Keith	MoF	250-749-6811
Brown, Patti	CFP	604-885-5905
Browne-Clayton, Shane	Riverside	250-762-3411
Carlson, Michael	MoF	250-260-4767
Cartwright, Charlie	MoF	250-387-6477
Cox, Keith	MoF	250-835-8626
Crowder, Tim	TimberWest	250-652-4211
Douglas, Diane	MoF	250-356-6721
Draper, Dale	MoF	250-387-3179
Ferguson, Craig	MoF	250-749-6811
Giampa, Gary	MoF	250-549-5576
Graham, Hilary	PRT	250-546-6713
Hak, Oldrich	Contractor	250-727-3989
Hollfreund, Clint	MoF	250-952-4127
Hooge, Bonnie	MoF	250-963-8416
Jaquish, Barry	MoF	250-260-4766
King, John	MoF	250-387-6476
Kolotelo, David	MoF	604-541-1683
Lee, Tim	VSOC	250-542-0833
Lohr, Carolyn	MoF	250-652-5600
Mehl, Helga	MoF	250-952-4172
Murphy, John	MoF	250-260-4767
Newton, Craig	Vizon Scitec Inc.	604-224-4331
Nicholson, George	Riverside	250-546-2293
Owens, John	UVic	250-652-4211
Painter, Roger	MoF	250-356-9276
Peiper, Greg	Riverside	250 546-2296
Phillips, Giselle	MoF	250 260 4756
Ponsford, Dave	MoF	250 387-1058
Rudolph, Dan	MoF	250-757-2015
Russell, John	MoF	250-749-6811
Stoehr, Michael	MoF	250-356-6209
Strong, Ward	MoF	250-549-5696
Van Neijenhuis, Annette	WFP	250-652-4023
Wagner, Rita	MoF	250 963-8416
Walsh, Chris	MoF	250-260-4777
Webber, Joe	MoF	250-537-8871
Wigmore, Bevin	TimberWest	250-652-4211
Woods, Jack	SelectSeed	250-748-9579

Back Cover

Grasshopper feeding on spruce shoots reduces the length of all needles equally, giving a brush-cut appearance.

Insect exclusion bag installed over cones of Lodgepole pine for *Leptoglossus* research.

Gall of Western Gall Rust (*Endocronartium harknesii*) on branch of lodgepole pine.



FGC

BRITISH COLUMBIA